

# Higher risk of heart defects in babies of mothers with type 1 diabetes

July 5 2018

Pregnant women with type 1 diabetes run a higher risk of having babies with heart defects, especially women with high blood glucose levels during early pregnancy, a study from Karolinska Institutet and the Sahlgrenska Academy in Sweden published in *The BMJ* shows.

It has long been known that patients with type 1 diabetes are at increased risk of complications. A new study now shows that pregnant [women](#) with type 1 diabetes are at a higher risk of having babies with heart defects.

"This confirms previous findings that there is a higher risk of birth defects, primarily of the heart," says Professor Jonas F. Ludvigsson at the Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, and consultant at the Paediatric Clinic at Örebro University Hospital. "The risk of birth defects is especially sensitive to factors during early pregnancy, and here blood glucose plays a vital part."

The study demonstrates a clear correlation between elevated levels of blood glucose (HbA1c) in the mother and the risk of heart defects in her baby. However, even those women who followed the current guidelines had a higher, albeit still small, risk of heart defects. The results show that 3.3 per cent of pregnant women with type 1 diabetes and [blood glucose levels](#) within the recommended span gave birth to a baby with a heart [defect](#). The corresponding figure for women without diabetes was 1.5 per cent.

Pregnant women with type 1 diabetes who had very [high blood glucose](#) levels (an HbA1c reading of 9.1 per cent or higher) were at much greater risk.

"Here, the risk of the baby having a heart defect was as much as 10.1 per cent—or one in every ten babies," says Professor Ludvigsson. "The reason why the risk of deformity can be linked to blood

glucose levels in early pregnancy is that it is then that the fetus's organs develop. Also, many women aren't aware that they're pregnant during the first few months."

This, he stresses, is why women must know about the dangers before trying to have children.

"There's an opportunity here for women to influence the risk of their baby developing a [heart](#) defect by keeping their blood glucose levels low. Yet we as doctors also know that many [pregnant women](#) struggle valiantly to keep their blood glucose down, as it is no easy task. The potential benefit of intensified insulin treatment to reduce the [risk](#) of [heart defects](#) should also be weighed against possible risks with hypoglycaemia in the mother and foetus," says Professor Ludvigsson.

The study was done by cross-referencing the National Diabetes Register with the National Patient Register and the Medical Birth Registry, and comparing 2,458 living newborns of mothers with type 1 diabetes with 1,159,865 babies of mothers without [diabetes](#). Since this is an observational study no definitive conclusions can be drawn regarding causality. The researchers are now planning to make further investigations in the field.

The study was financed with grants from the Swedish Diabetes Association, the Strategic Research Area in Epidemiology (SfoEpi) at Karolinska Institutet, the Swedish Research Council and Stockholm County Council.

**More information:** "Periconceptional glycemic control in type 1 diabetes and the risk of major birth defects: population based cohort study in Sweden", Jonas F. Ludvigsson, Martin Neovius, Jonas Söderling, Soffia Gudbjörnsdottir, Ann-Marie Svensson, Stefan Franzén, Olof Stephansson, Björn Pasternak. *BMJ*, online 5 July 2018.

Provided by Karolinska Institutet

APA citation: Higher risk of heart defects in babies of mothers with type 1 diabetes (2018, July 5)

retrieved 11 December 2022 from <https://medicalxpress.com/news/2018-07-higher-heart-defects-babies-mothers.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*