

Having hospital palliative care doesn't impact Tx intensity

September 27 2018



(HealthDay)—For adult patients in the intensive care unit (ICU), the



availability of hospital-based palliative care is not associated with inhospital treatment intensity, according to a study published in the September issue of the *Annals of the American Thoracic Society*.

May Hua, M.D., from the Columbia University College of Physicians and Surgeons in New York City, and colleagues conducted a retrospective study of adult ICU patients to examine the correlation between availability of hospital-based <u>palliative care</u> and treatment intensity. Data were included for 1,025,503 ICU patients in 151 hospitals.

The researchers found that 79.5 percent of the patients received care in a hospital with a palliative care program. Patients in hospitals with and without palliative care programs had a similar length of stay, with a statistically significant, but not clinically meaningful, difference (six versus six days; adjusted rate ratio, 1.04; 95 percent confidence interval, 1.03 to 1.05; P

"Currently, the measurable benefit of palliative care programs for critically ill <u>patients</u> may be the increased use of hospice facilities, as opposed to decreased health <u>care</u> use during an ICU-associated hospitalization," the authors write.

More information: Abstract/Full Text (subscription or payment may be required)

Editorial (subscription or payment may be required)

Copyright © 2018 HealthDay. All rights reserved.

Citation: Having hospital palliative care doesn't impact Tx intensity (2018, September 27) retrieved 30 April 2023 from

https://medicalxpress.com/news/2018-09-hospital-palliative-doesnt-impact-tx.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.