

Community health worker intervention beneficial

30 October 2018



quality of [care](#) (odds ratio, 1.8; 95 percent confidence interval, 1.4 to 2.4; P patients in the [intervention](#) group (odds ratio, 0.4; 95 percent confidence interval, 0.2 to 0.9).

"It's remarkable that these results were achieved not by an expensive drug or technology but from good old-fashioned social support," Kangovi said in a statement.

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)
[Editorial \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

(HealthDay)—A standardized community health worker (CHW)-delivered intervention, Individual Management for Patient-Centered Targets, improves patient-perceived quality of care and reduces hospitalizations for low-income patients with chronic diseases, according to a study published online Oct. 22 in *JAMA Internal Medicine*.

Shreya Kangovi, M.D., from the University of Pennsylvania in Philadelphia, and colleagues recruited 592 patients from three primary care facilities in a two-armed multicenter randomized clinical trial. Patients residing in a high-poverty ZIP code who were uninsured or publicly insured and had a diagnosis of two or more [chronic diseases](#) were recruited and randomly assigned to the CHW intervention or a control arm (goal setting only).

The researchers found similar improvements in self-rated physical health in the two arms (mean, 1.8 versus 1.6; P = 0.89). Intervention group patients had an increased likelihood of reporting the highest

APA citation: Community health worker intervention beneficial (2018, October 30) retrieved 28 July 2022 from <https://medicalxpress.com/news/2018-10-health-worker-intervention-beneficial.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.