

Over half of former ICU patients in the UK report symptoms of psychological disorders

22 November 2018

Patients in the UK who have survived critical illnesses requiring care in an intensive care unit (ICU) frequently report symptoms of anxiety, PTSD and/or depression, according to a study published in open access journal *Critical Care*. Those reporting symptoms of depression after critical illness appear to be at a greater risk of death.

Researchers at the University of Oxford investigated psychological disorders in a cohort of 4,943 of former ICU patients. They found that 46% of patients reported symptoms of anxiety, 40% reported symptoms of depression and 22% reported symptoms of PTSD, while 18% of patients in the study reported symptoms of all three psychological conditions.

Dr. Peter Watkinson, the corresponding [author](#) said: "Psychological problems after being treated for a critical illness in the ICU are very common and often complex when they occur. When symptoms of one psychological disorder are present, there is a 65% chance they will co-occur with symptoms of another psychological disorder."

To investigate possible links between treatment in an ICU and symptoms of psychological disorders, the authors asked a total of 4,943 patients who received treatment in one of 26 ICUs in the UK between 2006 and 2013, to complete a questionnaire on their symptoms of anxiety, depression and PTSD three months after discharge from ICU and again 12 months after discharge.

The authors found that patients who reported [symptoms of depression](#) were 47% more likely to die from any cause (all-cause mortality) during the first two years after discharge from the ICU than those who did not report these symptoms.

Dr. Peter Watkinson said: "Our findings suggest that [depression](#) following care of a critical illness in the ICU may be a marker of declining health and

clinicians should consider this when following up with former ICU patients."

The authors caution that the generalizability of the results outside of the UK may be limited as the data was only collected from UK based [patients](#). Furthermore, the observational nature of the study and its reliance of self-reported data mean that it does not allow for conclusions about cause and effect between ICU care and symptoms of [psychological disorders](#).

More information: Robert Hatch et al, Anxiety, Depression and Post Traumatic Stress Disorder after critical illness: a UK-wide prospective cohort study, *Critical Care* (2018). [DOI: 10.1186/s13054-018-2223-6](#)

Provided by BioMed Central

APA citation: Over half of former ICU patients in the UK report symptoms of psychological disorders (2018, November 22) retrieved 29 June 2022 from <https://medicalxpress.com/news/2018-11-icu-patients-uk-symptoms-psychological.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.