

Are most patients with fibromyalgia misdiagnosed?

6 February 2019

Recent studies have suggested that most people who have been diagnosed with fibromyalgia by physicians may not actually have the condition. A new *Arthritis Care & Research* study found that overall agreement between clinicians' diagnosis of fibromyalgia and diagnosis by published criteria is only fair.

For the study, 497 <u>patients</u> visiting a rheumatology clinic completed a health assessment questionnaire as well as a questionnaire that assessed fibromyalgia diagnostic variables used by the American College of Rheumatology. Patients were also evaluated and diagnosed by rheumatology clinicians.

Of the 497 patients, 121 (24.3 percent) satisfied fibromyalgia criteria while 104 (20.9 percent) received a clinician's diagnosis of fibromyalgia. The agreement between clinicians and criteria was 79.2 percent; however, agreement beyond chance was only fair. Physicians failed to identify 60 (49.6 percent) of criteria-positive patients and incorrectly identified 43 (11.4 percent) of criteria-negative patients.

"We have recently studied the same issue in 3,000 primary care patients and found about the same results. Therefore, our conclusions are secure," said lead author Dr. Frederick Wolfe, of the National Data Bank for Rheumatic Diseases.

In an accompanying editorial Dr. Don Goldenberg, of Oregon Health & Science University, noted that to conclude that expert physicians often misdiagnose fibromyalgia implies that published criteria are superior to expert clinical judgement for the <u>diagnosis</u> of individual patients. He stressed that this fails to account for many of the variables present in clinical encounters, including time spent in evaluating and categorizing each patient's multiple symptoms. "The diagnostic gold standard for <u>fibromyalgia</u> will continue to be the rheumatologist's expert opinion, not classification

criteria, no matter how well-refined and intentioned," he wrote. "This is the only way to capture the variability and severity of inter-related symptoms as they play out over time."

More information: *Arthritis Care & Research* (2019). DOI: 10.1002/acr.23731, onlinelibrary.wiley.com/journal/21514658

Provided by Wiley



APA citation: Are most patients with fibromyalgia misdiagnosed? (2019, February 6) retrieved 28 August 2022 from <u>https://medicalxpress.com/news/2019-02-patients-fibromyalgia-misdiagnosed.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.