

Personalized pain management

1 March 2019, by Niyati Vachharajani



These findings may assist in developing approaches in personalized pain medicine to predict opioid analgesic responses in patients.

Provided by Vanderbilt University

Credit: CC0 Public Domain

There is an urgent need to identify reliable predictors of opioid responses to develop personalized opioid therapy for chronic pain management. A goal is to target opioid therapy to patients most likely to benefit from treatment with few side effects and low risk of opioid misuse.

Stephen Bruehl, Ph.D.,and colleagues previously established an inverse correlation between levels of endogenous opioids (EOs), natural brain chemicals with pain-relieving properties, and analgesic responses to morphine.

Their latest study, published in the March issue of the journal *Pain*, tested whether circulating endocannabinoids, which stimulate the same brain receptors turned on by the <u>active ingredient</u> in marijuana, directly influence opioid analgesic responses or interact with EO function.

While they did not find that endocannabinoids directly affect morphine responses, they did find that the inverse association between EO activity and the analgesic and subjective effects of morphine was strongest in those with low endocannabinoid levels.

1/2



APA citation: Personalized pain management (2019, March 1) retrieved 31 May 2022 from https://medicalxpress.com/news/2019-03-personalized-pain.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.