


An antiviral gel may prevent genital herpes in women

1 March 2019, by Hannah Bae



Credit: CC0 Public Domain

In a paper recently published in the *Journal of Infectious Diseases*, researchers examined the effect of vaginal tenofovir 1 percent gel use on the risk of acquiring herpes simplex virus type 2, or HSV-2. The study was conducted through a secondary analysis of data from the VOICE study, the primary results of which were published in 2015.

According to the Centers for Disease Control and Prevention, more than one in six Americans ages 14 to 49 are infected with genital herpes. HSV-2 infection is the most common cause of genital herpes and also enhances the risk of acquiring and transmitting HIV. No vaccine currently exists for HSV-2.

"Women are especially susceptible to infection because it is more easily transmitted from an

infected man to his female sex partner than vice versa," said corresponding author Jeanne Marrazzo, M.D., director of the Division of Infectious Diseases at the University of Alabama at Birmingham. "A product that protects against HSV-2 could have an important public health impact."

Of the 566 HSV-2-negative participants, Marrazzo and her colleagues found that overall use of vaginal [tenofovir](#) 1 percent gel was associated with a 40 percent reduction of HSV-2 acquisition. Over the follow-up period, 92 new cases of HSV-2 occurred, with 77 in women with no plasma tenofovir detected and 15 with plasma tenofovir. Tenofovir measured in plasma was used as measure of gel use.

Marrazzo's study reanalyzed data collected in the [VOICE study](#), now specifically looking at the subgroup of women that adhered to daily gel usage and their rate of HSV-2 acquisition relative to women who did not adhere to gel usage.

In the original VOICE study, investigators studied HIV-1 pre-exposure prophylaxis in reproductive-age women in South Africa, Uganda and Zimbabwe. Participants were asked to regularly use study products, including vaginal tenofovir 1 percent gel, but adherence was low.

The recent study findings also align with the results of CAPRISA 004, a previous tenofovir gel study on HSV-2 acquisition reduction. The CAPRISA 004 study tested gel application strictly within 12 hours before and after vaginal intercourse.

"While both studies focus on the same product, our study is distinct in various ways, such as the emphasis on daily usage," Marrazzo said. "With our new findings, we can pay more attention to studying products combining tenofovir with other preventive materials in order to protect against HSV-2 and HIV—addressing both infections with multipurpose prevention products will maximize

protective impact for [women](#)."

More information: Jeanne M Marrazzo et al.
Tenofovir gel for prevention of herpes simplex virus
type-2 acquisition: findings from the VOICE Trial,
The Journal of Infectious Diseases (2019). [DOI:](#)
[10.1093/infdis/jiz045](https://doi.org/10.1093/infdis/jiz045)

Provided by University of Alabama at Birmingham

APA citation: An antiviral gel may prevent genital herpes in women (2019, March 1) retrieved 12
November 2022 from <https://medicalxpress.com/news/2019-03-antiviral-gel-genital-herpes-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.