

Younger female blood donors vulnerable to iron deficiency

March 21 2019



(HealthDay)—Blood donation is associated with iron deficiency among

both adolescent girls and younger adult women in the United States, according to a study recently published in *Transfusion*.

Eshan U. Patel, M.P.H., from the Johns Hopkins University School of Medicine in Baltimore, and colleagues used data from the 1999 to 2010 National Health and Nutrition Examination Survey to identify girls and women who reported their [blood donation](#) history in the preceding year and had serum ferritin (SF) measurements available. Participants were stratified by age (adolescents, 16 to 19 years: 2,419; adults, 20 to 49 years: 7,228).

The researchers found that in both adolescent and adult blood donors, geometric mean SF levels (ng/mL) were lower than in nondonors. Similarly, the prevalence of absent iron stores (SF

"These national data call for further development and implementation of blood donation practices aimed toward mitigating [iron deficiency](#)," the authors write.

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2019 [HealthDay](#). All rights reserved.

Citation: Younger female blood donors vulnerable to iron deficiency (2019, March 21) retrieved 4 February 2024 from <https://medicalxpress.com/news/2019-03-younger-female-blood-donors-vulnerable.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--