

Study finds young athletes are slightly more likely to have enlarged heart aortas

March 29 2019



Credit: CC0 Public Domain

Athletics training is associated with an increase in the athletes' heart ventricle wall thickness and cavity size. These changes are facilitated by the growth of heart muscle cells, or myocytes, in response to an

increased load on the heart from intensive physical exercise. Following episodes of 'detraining', the heart size returns to normal.

Two previous large studies involving athletes have shown that the thoracic [aorta](#) is also slightly increased in size; however, the significance of an enlarged aorta is unknown. Given that the aorta consists of a large amount of elastic tissue, it is possible that an enlarged aorta may represent reduced [elastic properties](#) and an inherent risk of aortic rupture. Although it might be expected that very tall athletes, such as [basketball players](#), would have a very dilated aorta based on their size, a recent American study showed that the aortic diameter rarely exceeds 40 mm.

A research team led by Professor Sanjay Sharma aimed to investigate the prevalence of an enlarged aorta in British athletes. They investigated 3781 athletes and 806 controls. Almost 2/3 of the athletes were male. The mean age in the group was 19 years old. Athletes trained for an average of over 16 hours per week.

The researchers found that athletes had a slightly larger aortic diameter compared with the control group; but this difference amounted to only 0.9 mm. The 99th percentile value for the aortic diameter in males and females was 40 mm and 38 mm respectively and this value was considered to represent an enlarged aortic diameter.

Five males and 6 females had an enlarged aortic diameter. The aortic diameter did not exceed 43 mm in any male or 41 mm in any female. These 11 athletes were followed up for over five years and underwent serial echocardiograms. They continued to exercise during that period. The researchers did not see any progressive enlargement in the aortic diameter in this group of athletes.

In conclusion, a small minority of athletes—approximately one in 300 or

0.3% - had a slightly enlarged aorta. The aortic diameter rarely exceeded 40 mm in males or 38 mm in females. During medium-term follow-up, there was no evidence of progressive enlargement of the aortic diameter in those athletes with an enlarged aorta.

Professor Sharma said: "Further long-term studies are required to understand the precise significance of an enlarged aortic diameter in athletes."

More information: Sabiha Gati et al. Prevalence and progression of aortic root dilatation in highly trained young athletes, *Heart* (2019). [DOI: 10.1136/heartjnl-2018-314288](https://doi.org/10.1136/heartjnl-2018-314288)

Provided by St. George's University of London

Citation: Study finds young athletes are slightly more likely to have enlarged heart aortas (2019, March 29) retrieved 11 February 2023 from <https://medicalxpress.com/news/2019-03-young-athletes-slightly-enlarged-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--