

More policy work needed to reduce use of smokeless tobacco, say researchers

April 3 2019



Chewing tobacco results in serious risk of cancer. Credit: University of York

Researchers at the University of York have shown that regulations on smokeless tobacco are still lacking, despite 181 countries agreeing to a common approach to controlling the demand and supply.

The study, published in *The Lancet Oncology*, highlighted that of the 181

countries using the WHO Framework Convention on Tobacco Control (FCTC), only 138 define smokeless tobacco in their statutes and 34 countries have so far reported levying tax on [smokeless tobacco products](#)

Just six countries check and regulate the content of smokeless tobacco products while only 41 mandate pictorial health warnings on these products.

Professor Kamran Siddiqi, from the University of York's Department of Health Sciences, said: "Smokeless tobacco is particularly popular in Asia and Africa and includes chewing tobacco as well as various types of nasal tobacco.

"They contain high levels of nicotine as well as cancer producing [toxic chemicals](#), making head and neck cancers common in those who consume smokeless tobacco products.

"Women are a particularly high-risk group, as cigarette smoking is less social acceptable in females in parts of Asia and Africa, resulting in smokeless tobacco being a popular alternative.

"We found that there is a policy implementation gap in smokeless [tobacco control](#), highlighting the need for increased global efforts to reduce the use of the products to catch-up with the progress made in curbing cigarette consumption."

Researchers, funded by the National Institute of Health Research (NIHR) are now working to establish a new global health group to address smokeless tobacco use in South Asia. The team will bring together researchers from around the world to critically assess policy and develop interventions to address the problems caused by [smokeless tobacco](#), particularly in India, Bangladesh and Pakistan.

Professor Ravi Mehrotra, Director of the Indian Council of Medical Research's National Institute of Cancer Prevention and Research (ICMR-NICPR), India, said: "Smokeless tobacco use is a public health concern in Southeast Asia, and beyond and requires a comprehensive approach to deal with the challenges identified in this study.

"The WHO FCTC Global Knowledge Hub on Smokeless Tobacco at ICMR-NICPR is committed to assisting in implementing the key recommendations in order to reduce the significant health burden."

More information: Ravi Mehrotra et al, Smokeless tobacco control in 180 countries across the globe: call to action for full implementation of WHO FCTC measures, *The Lancet Oncology* (2019). [DOI: 10.1016/S1470-2045\(19\)30084-1](https://doi.org/10.1016/S1470-2045(19)30084-1)

Provided by University of York

Citation: More policy work needed to reduce use of smokeless tobacco, say researchers (2019, April 3) retrieved 2 January 2024 from <https://medicalxpress.com/news/2019-04-policy-smokeless-tobacco.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.