


Low cholesterol linked to higher risk of bleeding stroke in women

10 April 2019



Credit: CC0 Public Domain

Lowering low-density lipoprotein (LDL) cholesterol reduces the risk of heart attacks and stroke, with an ideal value below 100 milligrams per deciliter (mg/dL). But can it be too low? A new study finds that women who have levels of LDL cholesterol 70 mg/dL or lower may be more than twice as likely to have a hemorrhagic stroke than women with LDL cholesterol levels from 100 to 130 mg/dL. The study is published in the April 10, 2019, online issue of *Neurology*, the medical journal of the American Academy of Neurology.

LDL [cholesterol](#) is called bad cholesterol because it can lead to fatty build-up in the arteries. The study also found that [women](#) with the lowest triglyceride levels, a fat found in the blood, had an increased risk of hemorrhagic stroke compared to those with the highest triglyceride levels.

Hemorrhagic strokes, also known as bleeding strokes, are much less common than ischemic strokes, when blood flow is blocked to the brain. They are also more difficult to treat and therefore more likely to be deadly.

"Strategies to lower cholesterol and triglyceride

levels, like modifying diet or taking statins, are widely used to prevent cardiovascular disease," said study author Pamela Rist, ScD, of Brigham and Women's Hospital in Boston and a member of the American Academy of Neurology. "But our large study shows that in women, very low levels may also carry some risks. Women already have a higher risk of stroke than men, in part because they live longer, so clearly defining ways to reduce their risk is important."

The study involved 27,937 women age 45 and older enrolled in the Women's Health Study who had total cholesterol, LDL cholesterol, high density lipoprotein (HDL) cholesterol, which is known as good cholesterol, and triglycerides measured at the start of the study. Researchers then reviewed medical records to determine how many women had a bleeding stroke.

With an average of 19 years of follow-up, researchers identified 137 women who had a bleeding stroke.

Nine out of 1,069 women with cholesterol 70 mg/dL or lower, or 0.8 percent, had a bleeding stroke, compared to 40 out of 10,067 women with cholesterol 100 mg/dL up to 130 mg/dL, or 0.4 percent. After adjusting for other factors that could affect risk of stroke, such as age, smoking status, high blood pressure and treatment with cholesterol-lowering medications, researchers found that those with very low LDL cholesterol were 2.2 times more likely to have a bleeding stroke.

For triglyceride levels, researchers divided the women into four groups. Women in the group with the lowest levels had fasting levels 74 mg/dL or lower, or non-fasting levels of 85 mg/dL or lower. Women in the group with the highest levels had fasting levels that were higher than 156 mg/dL, or non-fasting levels that were higher than 188 mg/dL.

Researchers found that 34 women of the 5,714

women with the lowest levels of triglycerides, or 0.6 percent, had a bleeding stroke, compared to 29 women of the 7,989 with the highest triglycerides, or 0.4 percent. After adjusting for other factors that could affect risk, researchers found that those with the lowest triglycerides were twice as likely to have a bleeding stroke.

No difference in risk was found for total cholesterol or HDL cholesterol.

"Women with very low LDL cholesterol or low triglycerides should be monitored by their doctors for other stroke risk factors that can be modified, like [high blood pressure](#) and smoking, in order to reduce their risk of hemorrhagic stroke," Rist said. "Also, additional research is needed to determine how to lower the risk of [hemorrhagic stroke](#) in women with very low LDL and low triglycerides."

One limitation of the study was that cholesterol and triglyceride levels were only measured once at the beginning of the study. Also, a large number of the women had already reached menopause when those levels were measured, which prevented researchers from examining whether menopause status may influence the link between cholesterol and triglyceride levels and bleeding [stroke](#).

Provided by American Academy of Neurology

APA citation: Low cholesterol linked to higher risk of bleeding stroke in women (2019, April 10) retrieved 24 October 2022 from <https://medicalxpress.com/news/2019-04-cholesterol-linked-higher-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.