

# Thiazide diuretics reduce risk of fractures in people with Alzheimer's disease

23 April 2019



Long-term use of thiazides reduced the risk of low energy fractures in people with Alzheimer's disease. Credit: Raija Törrönen

The use of thiazide diuretics was associated with a decreased risk of low energy fractures in people with Alzheimer's disease, a new study from the University of Eastern Finland shows. The association was found in long-term use exceeding three years; however, shorter term use did not reduce the risk of fractures. Thiazides are typically

prescribed to treat hypertension. The results were published in *Osteoporosis International*.

Hip fracture was the most common low energy fracture in the study population, and the risk of this fracture type decreased by 30% in long-term thiazide use. The association may be related to the fact that thiazides decrease urinary calcium excretion and increase bone mineral density, which may decrease the risk of fracture in case of a fall. Persons with Alzheimer's disease have an increased risk of falling, as the disease leads to problems in balance and spatial orientation.

The association of thiazide use and a reduced risk of low energy [fractures](#) disappeared approximately one month after use was discontinued. Thiazides are used in the treatment of hypertension, and in this study, thiazide use was taken into account also when these drugs were combined with other antihypertensive drugs in the same tablet. Thiazides cannot be recommended for the prevention of fractures alone, as their use is known to be associated with postural hypotension and consequent falls.

The study was based on the nationwide register-based MEDALZ study conducted at the University of Eastern Finland. For this study, 10,416 community-dwelling persons diagnosed with Alzheimer's disease who had a low energy fracture were included and compared with 31,099 matched controls without a fracture.

**More information:** H. Taipale et al. Long-term thiazide use and risk of low-energy fractures among persons with Alzheimer's disease—nested case-control study, *Osteoporosis International* (2019). [DOI: 10.1007/s00198-019-04957-0](https://doi.org/10.1007/s00198-019-04957-0)

Provided by University of Eastern Finland

APA citation: Thiazide diuretics reduce risk of fractures in people with Alzheimer's disease (2019, April 23) retrieved 11 September 2022 from <https://medicalxpress.com/news/2019-04-thiazide-diuretics-fractures-people-alzheimer.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*