

Dietary cholesterol and egg consumption do not increase the risk of stroke

May 20 2019



Credit: CC0 Public Domain

A new study from the University of Eastern Finland shows that a moderately high intake of dietary cholesterol or consumption of up to one egg per day is not associated with an elevated risk of stroke.

Furthermore, no association was found in carriers of the APOE4 phenotype, which affects cholesterol metabolism and is remarkably common among the Finnish population. The findings were published in the *American Journal of Clinical Nutrition*.

Findings from earlier studies addressing the association of dietary [cholesterol](#) or egg intake with the risk of stroke have been contradictory. Some studies have found an association between high dietary cholesterol intake and an increased risk of stroke, while others have associated the consumption of eggs, which are high in cholesterol, with a reduced risk of stroke. For most people, dietary cholesterol plays a very small role in affecting their serum cholesterol levels. However, in carriers of the apolipoprotein E phenotype 4—which significantly impacts [cholesterol metabolism](#)—the effect of dietary cholesterol on serum cholesterol levels is greater. In Finland, the prevalence of APOE4, which is a hereditary variant, is exceptionally high, with approximately one third of the population presenting as carriers. Yet, [research data](#) on the association between a high intake of dietary cholesterol and the risk of stroke in this population group has not been available until now.

The [dietary habits](#) of 1,950 men aged between 42 and 60 years with no baseline diagnosis of a [cardiovascular disease](#) were assessed at the onset the Kuopio Ischaemic Heart Disease Risk Factor Study, KIHDS, in 1984–1989 at the University of Eastern Finland. APOE phenotype data were available for 1,015 of the men participating in the study. Of those, 32 percent were known carriers of APOE4.

During a follow-up of 21 years, 217 men were diagnosed with stroke. The study found that neither dietary cholesterol nor egg consumption was associated with the risk of stroke – not even in carriers of APOE4.

The findings suggest that moderate cholesterol intake or daily egg consumption are not associated with the risk of [stroke](#), even in persons

who are genetically predisposed to a greater effect of dietary cholesterol on serum cholesterol levels. In the highest control group, the [study participants](#) had an average daily dietary cholesterol intake of 520 mg and they consumed an average of one egg per day, which means that the findings cannot be generalised beyond these levels. One egg contains approximately 200 mg of cholesterol. In this study, about a fourth of the total dietary cholesterol consumed came from eggs.

Furthermore, the generalisability of this study is also weakened by the fact that the study population did not have a pre-existing cardiovascular disease at baseline and the size of the study population was relatively small. Therefore, the findings of the study should be verified in a larger cohort as well as in people with a pre-existing cardiovascular disease, who are currently advised to limit their intake of cholesterol and [eggs](#).

More information: Anna M Abdollahi et al. Egg consumption, cholesterol intake, and risk of incident stroke in men: the Kuopio Ischaemic Heart Disease Risk Factor Study, *The American Journal of Clinical Nutrition* (2019). [DOI: 10.1093/ajcn/nqz066](https://doi.org/10.1093/ajcn/nqz066)

Provided by University of Eastern Finland

Citation: Dietary cholesterol and egg consumption do not increase the risk of stroke (2019, May 20) retrieved 4 February 2024 from <https://medicalxpress.com/news/2019-05-dietary-cholesterol-egg-consumption.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--