

# Body parts respond to day and night independently from brain, studies show

30 May 2019



"The future implications of our findings are vast," says Paolo Sassone-Corsi, senior author of one of the two studies on circadian clocks published today in the journal *Cell*. He directs UCI's Center for Epigenetics and Metabolism and is a Donald Bren Professor of Biological Chemistry. Penny Lee / UCI School of Medicine

Can your liver sense when you're staring at a television screen or cellphone late at night? Apparently so, and when such activity is detected, the organ can throw your circadian rhythms out of whack, leaving you more susceptible to health problems.

That's one of the takeaways from two new studies by University of California, Irvine scientists working in collaboration with the Institute for Research in Biomedicine in Barcelona, Spain.

The studies, published today in the journal *Cell*, used specially bred [mice](#) to analyze the network of internal clocks that regulate metabolism. Although researchers had suspected that the body's various circadian clocks could operate independently from the central clock in the hypothalamus of the brain, there was previously no way to test the theory, said Paolo Sassone-Corsi, director of UCI's Center for

Epigenetics and Metabolism and senior author of one of the studies.

To overcome that obstacle, scientists figured out how to disable the entire circadian system of the mice, then jump-start individual clocks. For the experiments reported in the *Cell* papers, they activated clocks inside the liver or skin.

"The results were quite surprising," said Sassone-Corsi, Donald Bren Professor of Biological Chemistry. "No one realized that the liver or skin could be so directly affected by light."

For example, despite the shutdown of all other body clocks, including the central brain clock, the liver knew what time it was, responded to light changes as day shifted to night and maintained critical functions, such as preparing to digest food at mealtime and converting glucose to energy.

Somehow, the liver's [circadian clock](#) was able to detect light, presumably via signals from other organs. Only when the mice were subjected to constant darkness did the liver's clock stop functioning.

In upcoming studies, UCI and Barcelona researchers will phase in other internal clocks to see how different organs communicate with each other, Sassone-Corsi said.

"The future implications of our findings are vast," he noted. "With these mice, we can now begin deciphering the metabolic pathways that control our [circadian rhythms](#), aging processes and general well-being."

In earlier studies, Sassone-Corsi has examined how circadian clocks can be rewired by such factors as sleep deprivation, diet and exercise. Exposure to computer, television or cellphone light just before bed can also scramble internal clocks.

Because of modern lifestyles, it's easy for people's circadian systems to get confused, he said. In turn, that can lead to depression, allergies, premature aging, cancer and other health problems. Further mice experiments could uncover ways to make human internal clocks "less misaligned," Sassone-Corsi added.

**More information:** Patrick-Simon Welz et al, BMAL1-Driven Tissue Clocks Respond Independently to Light to Maintain Homeostasis, *Cell* (2019). [DOI: 10.1016/j.cell.2019.05.009](https://doi.org/10.1016/j.cell.2019.05.009)

Provided by University of California, Irvine

APA citation: Body parts respond to day and night independently from brain, studies show (2019, May 30) retrieved 31 October 2022 from <https://medicalxpress.com/news/2019-05-body-day-night-independently-brain.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*