

Formation of habitual use drives cannabis addiction

13 June 2019



Credit: CC0 Public Domain

A shift from brain systems controlling reward-driven use to habit-driven use differentiates heavy cannabis users who are addicted to the drug from users who aren't, according to a study in *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*. The findings help explain how the brain becomes dependent on cannabis, and why not all cannabis users develop an addiction, even with long-term regular use.

In the study, researchers at the University of Electronic Science and Technology of China and University of Bonn, Germany, used brain imaging to monitor [neural activity](#) when cannabis users viewed images associated with cannabis use, referred to as drug cues. Although all of the cannabis users in the study reported heavy use, only some were dependent on the drug. Both dependent and non-dependent [cannabis users](#) had exaggerated responses in a brain region that processes reward—the [ventral striatum](#)—compared with people who didn't use cannabis. Interestingly, the dependent users also had larger responses in a brain region that forms habits—the dorsal striatum.

"The present findings reflect that heavy cannabis use is promoted by changes in the brain's reward system—however, these changes alone may not fully explain addictive use. Addictive use may rather be driven by changes in [brain systems](#) that promote habitual—automatic—use, which also may explain the fact that addicts continue use despite a lack of experiencing rewarding effects of the drug. As such, their behavior has become under the control of the drug cues, rather than the actual reward expectation," said lead author Benjamin Becker, Ph.D.

Dependent users also had increased responses in other regions throughout the brain, including regions that attribute importance to things, for example, drug cues. This suggests that the development of cannabis addiction incorporates additional [brain regions](#) that may strengthen a person's desire to seek the drug.

"Cannabis is now legal for medical and recreational use in many parts of the United States and the health impacts of this development are still being understood," said Cameron Carter, MD, Editor of *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*. "These findings are important insights that can help us better understand why some individuals might be more likely to become addicted to cannabis," he added.

Differentiating the unique brain circuits behind dependent cannabis use could also be useful for understanding how to combat the problem of [cannabis](#) addiction. "The identification of the [dorsal striatum](#) and habitual behavior as a driver of addiction may allow the development of more specific treatment approaches to increase treatment success," said first author Xinqi Zhou.

More information: Xinqi Zhou et al, Cue-reactivity in the ventral striatum characterizes heavy cannabis use, whereas reactivity in the dorsal striatum mediates dependent use, *Biological*

*Psychiatry: Cognitive Neuroscience and
Neuroimaging* (2019). DOI:
[10.1016/j.bpsc.2019.04.006](https://doi.org/10.1016/j.bpsc.2019.04.006)

Provided by Elsevier

APA citation: Formation of habitual use drives cannabis addiction (2019, June 13) retrieved 13 June 2021 from <https://medicalxpress.com/news/2019-06-formation-habitual-cannabis-addiction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.