

# Poor sleep quality and fatigue plague women with premature ovarian insufficiency

16 July 2019



Credit: CC0 Public Domain

Sleep disturbances are a frequent complaint of women in the menopause transition and postmenopause. A new study demonstrates that women with premature ovarian insufficiency (POI) who are receiving hormone therapy have poorer sleep quality and greater fatigue than women of the same age with preserved ovarian function. Study results are published online in *Menopause, the journal of The North American Menopause Society (NAMS)*.

It is estimated that 40% to 50% of menopausal and [postmenopausal women](#) complain about [sleep problems](#). Sleep problems include difficulty falling asleep and/or staying asleep, as well as waking up too early. Complicating matters is the fact that women with insomnia generally complain more about body pain, headaches, daytime dysfunction, mood disorders, fatigue, and decreased work productivity. Although some of the problems are related to other common symptoms of menopause such as hot flashes, not all sleep issues can be traced back to these root causes.

Although [numerous studies](#) have been conducted

about the sleep patterns of menopausal and postmenopausal women in general, this newest study from Brazil is thought to be the first to specifically evaluate the sleep quality in women with POI. This condition differs from premature menopause because women with POI can still have irregular or occasional periods for years and might even become pregnant. The Brazilian study found that women with POI who are receiving hormone therapy have poorer sleep quality, largely as a result of taking longer to fall asleep. These women were also found to have a higher fatigue index and were more likely to use sleep-inducing medications compared with comparably aged women who still had full ovarian function.

Study results appear in the article "Sleep quality and fatigue in women with premature ovarian insufficiency receiving hormone therapy: a [comparative study](#)."

"This study shows that women with POI have poor sleep quality despite the use of [hormone therapy](#). Another interesting finding from the study is that total sleep quality in women with POI was directly related to the number of children they had and overall was similar to sleep quality in women without POI. This speaks to the scope of the problem when it comes to [sleep disturbances](#) and the important and often under-recognized factors that contribute to sleep complaints being more common in women than in men," says Dr. Stephanie Faubion, NAMS medical director.

**More information:** Cristina Laguna Benetti-Pinto et al, Sleep quality and fatigue in women with premature ovarian insufficiency receiving hormone therapy, *Menopause* (2019). DOI: [10.1097/GME.0000000000001379](https://doi.org/10.1097/GME.0000000000001379)

Provided by The North American Menopause Society

APA citation: Poor sleep quality and fatigue plague women with premature ovarian insufficiency (2019, July 16) retrieved 25 June 2022 from <https://medicalxpress.com/news/2019-07-poor-quality-fatigue-plague-women.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*