

Statistical analyses reveal the shape of epileptic seizures

July 23 2019



Yuan Wang (left) and Hernando Ombao discuss their statistical models. Credit: KAUST

Pyramidal graphs resulting from statistical analyses of EEG recordings can improve our understanding of epileptic seizures.

A statistical approach squeezes more detailed information out of a current method of measuring [brain signals](#) in [epileptic seizures](#), adding new insight into how these signals originate and spread.

Visual inspection of electroencephalography (EEG) recordings of epilepsy patients before and during a seizure is a fairly effective method for detecting the part of the [brain](#) that can benefit from surgical treatment. But it's not sufficient for more challenging cases.

Now, an approach developed by KAUST biostatistician Hernando Ombao and colleagues Yuan Wang of the University of Wisconsin–Madison and Moo K. Chung of the University of California, Irvine, digs deeper into the features of an EEG and can detect abnormalities in [brain regions](#) even before a seizure takes place.

The method could provide a clinically useful tool for seizure localization, according to Chung and

Wang.

The approach stems from a field of mathematics that analyzes large and complex datasets by studying shape representations of the data and its interactions. Analyzing these shapes provides information on patterns that exist within the data. The team applied their method, known as a topological data analysis framework, to see what they could learn from an EEG recording conducted before and during an epileptic seizure.

The statistical approach removes noise from the EEG recording, providing cleaner signals. A series of shapes are then drawn that directly relate to the signals in the recordings. The final pyramidal shapes (persistence landscapes) that represent the signals coming from each electrode placed on the scalp provide a good picture of where the seizure originates in the brain and how it spreads.

The analysis of the patient's EEG recording showed that the seizure originated from a region in or around the electrodes measuring signals from the left temporal lobe of the brain. It then spread to the right temporal lobe.

Further simulation studies showed that the test was robust and sensitive, even when the signal was buried under noise.

"Epileptologists should enhance their toolboxes of data analysis by adding methods like this one that capture topological features as part of their assessment of [seizure](#) foci in more challenging cases of epilepsy," says Ombao.

The team next plans to test their framework on large samples of EEG recordings to clinically validate their findings. Ombao is also developing [statistical methods](#) to study the impact of shocks to the brain, such as in epilepsy or stroke, on the communication network between brain regions and nerve cell populations.

More information: Yuan Wang et al. Topological data analysis of single-trial electroencephalographic signals, *The Annals of Applied Statistics* (2018). DOI: [10.1214/17-AOAS1119](https://doi.org/10.1214/17-AOAS1119)

Provided by King Abdullah University of Science and Technology

APA citation: Statistical analyses reveal the shape of epileptic seizures (2019, July 23) retrieved 8 December 2022 from <https://medicalxpress.com/news/2019-07-statistical-analyses-reveal-epileptic-seizures.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.