

# Oral semaglutide reduces HbA1c, weight in patients with T2DM

July 31 2019

---



(HealthDay)—Compared with placebo, oral semaglutide monotherapy is

associated with superior and clinically relevant improvements in glycated hemoglobin (HbA1c) and weight loss among patients with type 2 diabetes, according to a study published online July 18 in *Diabetes Care*.

In a phase 3a, randomized, double-blind, placebo-controlled, parallel-group trial, Vanita R. Aroda, M.D., from Brigham and Women's Hospital in Boston, and colleagues evaluated the safety and efficacy of oral semaglutide, the first oral glucagon-like peptide 1 receptor agonist, as monotherapy in patients with type 2 diabetes managed by diet and exercise alone. The authors randomly assigned (1:1:1:1) 703 patients at 93 international sites to either once-daily oral semaglutide 3 mg, 7 mg, 14 mg, or placebo.

Over 26 weeks, the researchers found that oral semaglutide reduced HbA1c as well as body weight. The most common adverse events associated with oral semaglutide included mild-to-moderate transient gastrointestinal events. Discontinuation of the trial product occurred in 2.3 to 7.4 percent taking oral semaglutide and 2.2 percent among those taking placebo.

"Ongoing additional studies in the PIONEER [Peptide Innovation for Early Diabetes Treatment] program will further define its effect when used in combination with other glucose-lowering therapies and in other populations (e.g., in those with high cardiovascular risk or [renal impairment](#)) of interest," the authors write.

Several authors disclosed financial ties to [pharmaceutical companies](#), including Novo Nordisk, which manufactures semaglutide and funded the study.

**More information:** [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2019 [HealthDay](#). All rights reserved.

Citation: Oral semaglutide reduces HbA1c, weight in patients with T2DM (2019, July 31)  
retrieved 5 October 2023 from

<https://medicalxpress.com/news/2019-07-oral-semaglutide-hba1c-weight-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.