


Low vitamin D levels linked to non-motor symptoms in patients with Parkinson's disease

7 August 2019



the Second Affiliated Hospital of Soochow University, in China.

More information: Hui?Jun Zhang et al, Relationship between 25?Hydroxyvitamin D, bone density, and Parkinson's disease symptoms, *Acta Neurologica Scandinavica* (2019). [DOI: 10.1111/ane.13141](https://doi.org/10.1111/ane.13141)

Provided by Wiley

Immunohistochemistry for alpha-synuclein showing positive staining (brown) of an intraneural Lewy-body in the Substantia nigra in Parkinson's disease. Credit: Wikipedia

In an *Acta Neurologica Scandinavica* study of 182 patients with Parkinson's disease and 185 healthy controls, patients with Parkinson's disease had significantly lower levels of vitamin D in their blood. Also, patients with lower vitamin D levels were more likely to fall, and to experience sleep problems, depression, and anxiety.

The findings suggest that vitamin D supplementation may help to treat non-motor symptoms associated with Parkinson's disease.

"As various non-[motor symptoms](#) place a burden on individuals with Parkinson's [disease](#) and their caregivers, vitamin D might be a potential add-on therapy for improving these neglected symptoms," said senior author Chun Feng Liu, MD, Ph.D., of

APA citation: Low vitamin D levels linked to non-motor symptoms in patients with Parkinson's disease (2019, August 7) retrieved 11 August 2022 from <https://medicalxpress.com/news/2019-08-vitamin-d-linked-non-motor-symptoms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.