

# Weight change and bone health in older adults with obesity

September 5 2019

---

Weight loss in older adults is accompanied by loss in bone mineral density (BMD) and an increased risk of bone fracture.

A new study published in *Obesity* found that loss of hip BMD persists in the year following a weight loss intervention among [older adults](#) with obesity, regardless of whether they regain weight.

The study also found that losses in fat mass and lean mass contribute to reduced hip BMD; yet, loss in fat mass may signal improved trabecular bone score, which is a measure of bone texture and is a marker for the risk of osteoporosis.

The findings emphasize the importance of identifying and implementing interventions that can target [fat mass](#), but not lean mass, loss to optimize bone health during [weight loss](#) in older adults.

**More information:** Daniel E. Kammire et al, Effect of Weight Change Following Intentional Weight Loss on Bone Health in Older Adults with Obesity, *Obesity* (2019). [DOI: 10.1002/oby.22604](https://doi.org/10.1002/oby.22604)

Provided by Wiley

Citation: Weight change and bone health in older adults with obesity (2019, September 5) retrieved 25 February 2023 from <https://medicalxpress.com/news/2019-09-weight-bone-health->

[older-adults.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.