

# Low body-mass index with abdominal obesity is associated with worse heart failure outcomes in Asian

24 September 2019

---



Having a lower body-mass index (BMI), but also with having a higher waist-to-height ratio (WHtR), is associated with worse outcomes among Asian patients with heart failure, according to a study published September 24 in the open-access journal *PLOS Medicine* by Carolyn Lam of the National Heart Centre Singapore, and colleagues. Credit: Pexels, Pixabay

Having a lower body-mass index (BMI), but also having a higher waist-to-height ratio (WHtR), is associated with worse outcomes among Asian patients with heart failure, according to a study published September 24 in the open-access journal *PLOS Medicine* by Carolyn Lam of the National Heart Centre Singapore, and colleagues. As noted by the authors, the combined use of BMI and abdominal measures could potentially inform heart failure management better, especially among the particularly vulnerable patients with low BMI and high WHtR in Asia.

Among [heart failure patients](#), Asian patients are more likely to be lean (have a low BMI,

APA citation: Low body-mass index with abdominal obesity is associated with worse heart failure outcomes in Asian (2019, September 24) retrieved 17 September 2022 from <https://medicalxpress.com/news/2019-09-body-mass-index-abdominal-obesity-worse.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*