


Risk of heart valve infections rising in hospitals

29 September 2019



Credit: CC0 Public Domain

People with heart disease or defective or artificial heart valves are at increased risk of developing a potentially deadly valve infection. Rutgers researchers reported that new risk factors for this condition have emerged and that an increasing number of patients admitted to hospitals for other diseases are at risk of contracting this potentially lethal cardiac infection.

The study, which was published online prior to print in *The American Journal of Cardiology*, highlights the need for hospitals to develop ways to prevent this serious [infection](#) in the [heart](#).

The American Heart Association had recommended that all people at risk for heart valve infections ([infective endocarditis](#))—typically caused by bacteria entering the bloodstream through the mouth, gastrointestinal or genitourinary tract—take antibiotics. In 2007, the guidelines were revised to recommend antibiotics only for those determined to be at high risk for infection.

"In the past, infective endocarditis was associated with [rheumatic heart disease](#) and most often caused by bacteria in the mouth," said lead author

Abel Moreyra, a Professor of Medicine at Rutgers Robert Wood Johnson Medical School. "However, new risk factors, such as intravenous opiate abuse, compromised immune systems, hemodialysis and implanted heart devices have emerged."

To understand how guideline changes affected the rate of infections, the researchers analyzed 21,443 records of people who were diagnosed with infective endocarditis in New Jersey hospitals from 1994 to 2015. They made a startling discovery: Beginning in 2004 and continuing thereafter, there was a significant decline in the number of patients hospitalized with infective endocarditis as the primary diagnosis for their reason for admission and a significant increase in the number of patients developing the infection in the hospital, or a secondary diagnosis. In total, 9,191 people were hospitalized with infective endocarditis as the primary diagnosis and 12,252 with secondary diagnosis.

Moreyra attributes the decline in primary diagnosis to improved [dental care](#) and the rarity today of rheumatic [heart disease](#), where streptococcus plays a predominant role in the infection. "However, 60 percent of infective endocarditis that developed after admission were caused by a different microorganism, staphylococcus bacteria, which is abundant in hospitals and implicates health care as a possible source of infection," he said.

This important analysis of the different time trends of primary and secondary diagnosis of infective endocarditis can help hospitals tailor different strategies for the prevention of this potentially lethal infection, Moreyra said.

More information: Abel E. Moreyra et al, Trends in Hospitalization for Infective Endocarditis as a Reason for Admission or a Secondary Diagnosis, *The American Journal of Cardiology* (2019). [DOI: 10.1016/j.amjcard.2019.04.045](#)

Provided by Rutgers University

APA citation: Risk of heart valve infections rising in hospitals (2019, September 29) retrieved 6 November 2022 from <https://medicalxpress.com/news/2019-09-heart-valve-infections-hospitals.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.