

What patients expect influences pain management

8 October 2019

An article in the current issue of *Psychotherapy and Psychosomatics* explores the psychological aspects of pain management with particular reference to placebo infusions in back pain.

Increasing evidence for the efficacy of analgesic placebo effects in [laboratory studies](#) with healthy persons raises the question of whether placebos could be used to improve the treatment of [pain](#) patients. Expectancies play a central role in shaping the analgesic placebo but also nocebo effects. This study investigated to what extent a sham opioid infusion (saline solution) produces sustained clinically relevant placebo and nocebo effects in chronic back pain patients.

Fifty-nine patients received the sham opioid infusion applied via a large drain dressing and were compared to 14 control patients without intervention while experimental pain stimuli were applied. All subjects were told that the infusion would decrease pain although in rare cases pain increase would be possible (induction of [expectancy](#)). In addition, conditioning was introduced where the participants either experienced a decrease in experimental pain (n= 17; placebo conditioning), an increase (n= 21; nocebo conditioning), or no change (n= 21, no conditioning).

Results showed that compared to the [control group](#), all infusion groups showed positive treatment expectancies and significantly reduced clinical back pain (primary outcome) and pain-related disability (secondary outcome, assessed by self-reported functional capacity and perceived impairment of mobility). Even the nocebo conditioned group experiencing increased experimental pain developed positive treatment expectancies followed by reduced pain experience. Positive treatment expectancies and relief in clinical [back pain](#) were significantly positively correlated.

The authors concluded that these findings suggest that it may be beneficial to explicitly shape and integrate treatment expectancies into clinical pain management.

More information: Julia Schmitz et al. Positive Treatment Expectancies Reduce Clinical Pain and Perceived Limitations in Movement Ability Despite Increased Experimental Pain: A Randomized Controlled Trial on Sham Opioid Infusion in Patients with Chronic Back Pain, *Psychotherapy and Psychosomatics* (2019). [DOI: 10.1159/000501385](#)

Provided by Journal of Psychotherapy and Psychosomatics

APA citation: What patients expect influences pain management (2019, October 8) retrieved 17 June 2022 from <https://medicalxpress.com/news/2019-10-patients-pain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.