

Expectant mothers' pregnancy-related anxiety may alter how infant brains respond to sad speech

November 12 2019



The brain scanning equipment uses fibre optics held to the baby's head with a gauze. Credit: University of Turku & Aalto University

A study has shown a potential link between pregnancy-related anxiety

and how a baby's brains respond to sad speech. Researchers at Aalto University and the University of Turku in Finland showed that mothers with high anxiety scores at 24 weeks of pregnancy gave birth to babies who had reduced brain responses to sad-sounding speech. The effect was significantly smaller at 34 weeks of pregnancy, suggesting the effects of pregnancy-related anxiety may be transferred more easily to the unborn baby in mid- rather than late pregnancy. Studying a larger group would make it possible to understand the behavioral implications of the observed changes.

"Areas of the baby's brain that deal with emotion and [speech](#) were less active when listening to sad speech if the baby's mothers had reported high pregnancy-related anxiety" said Dr. Ilkka Nissilä, a research fellow at Aalto University who is one of the authors of the study.

Pregnancy-related anxiety refers to worries specifically concerning the pregnancy, such as worries about changes in appearance, labor and birth, the health of the developing child and future parenting. Previous studies have already shown a link between anxiety during [pregnancy](#) and neurodevelopment in infants, which prompted the researchers to investigate emotional speech and [anxiety](#) in this study.

"While we can observe a change in brain response, we can't say for sure how it is related to behavior, or how the changes we've observed affect the child over time " said Dr. Nissilä. "What we know for certain is that a larger study with more subjects is needed to understand how such observations made using neuroimaging are related to the development of the babies as they grow up."

Professor Hasse Karlsson, professor of integrative neuroscience and psychiatry at the University of Turku said "One of the advantages of the FinnBrain study is that the babies will be followed up for several years. This makes it possible to later find out if this finding is related to any

clinically relevant outcomes."

The study examined the data of 19 mother-baby pairs from Finland. The brain scanning was done with a technique called diffuse optical tomography or DOT, which uses a set of light sources and detectors attached on the side of the baby's head to measure changes in blood flow in the [brain](#). The method allows the study to take place while the baby is in its mother's lap, and is almost silent, as opposed to the fMRI scans often used in functional neuroscience experiments which require lying still in a loud scanner. The DOT equipment meant that the researchers could easily play speech sounds to the [babies](#) whilst simultaneously scanning their brains. They hope to be able to improve the DOT technology further for future work by making it wireless.

The work was published in the *Journal of Affective Disorders*.

More information: Ambika Maria et al. Relationship between maternal pregnancy-related anxiety and infant brain responses to emotional speech – a pilot study, *Journal of Affective Disorders* (2019). [DOI: 10.1016/j.jad.2019.10.047](https://doi.org/10.1016/j.jad.2019.10.047)

Provided by Aalto University

Citation: Expectant mothers' pregnancy-related anxiety may alter how infant brains respond to sad speech (2019, November 12) retrieved 15 February 2024 from <https://medicalxpress.com/news/2019-11-mothers-pregnancy-related-anxiety-infant-brains.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--