


New study identifies potential path forward for brachial plexus injury recovery

January 16 2020, by Brandi Wampler



Cody Smith chats with a graduate student in his lab. Credit: Matt Cashore/University of Notre Dame

On average, an estimated three out of every 1,000 newborns will suffer a brachial plexus injury during birth, damaging the bundle of nerves that connect the brain and spinal cord to the shoulders, arms and hands. In the most traumatic cases, even with surgery and physical therapy as an infant, there is no treatment that can guarantee a full recovery.

However, a new study from the University of Notre Dame has identified a strategy that may support the regeneration of nerves affected by a brachial plexus [injury](#). The findings show that there could be a new path forward for a full behavioral recovery.

"Early on, our model showed that the [nerve](#) regeneration process after a brachial plexus injury differed from how these nerves connect the peripheral and central nervous systems during early development," said Cody Smith, the Elizabeth and Michael Gallagher Assistant Professor of Biological Sciences and co-author of the study. "My lab worked to recreate that development-like process by using a chemotherapy treatment, paclitaxel, which helped stabilize axon invasion for spinal cord re-entry."

During [early development](#), nerves known as dorsal root ganglia (DRG) sensory axons enter the spinal cord to then connect with specific neurons, which provide motor and sensory function for hands, arms and shoulders. However, after a brachial plexus injury, these DRG sensory axons are unable to penetrate the spinal cord barrier when attempting to regenerate. For the study, which was published in *Cell Reports*, the Notre Dame researchers aimed to identify a process that would allow severed DRG sensory axons to enter the spinal cord when regenerating after an injury within their model.

Once the DRG sensory axons penetrated the [spinal cord](#), the researchers were able to establish connection between the nerves and with neuronal "partners," allowing for full functional recovery in the model after 48 hours. Although the study was intended to mimic regeneration after obstetrical brachial plexus injuries, there is potential for it to benefit those who suffer a [brachial plexus injury](#) later in life as a result of an accident or other trauma.

"Moving forward, our goal is to use our studies during development to

identify more specific strategies for regeneration," said Smith. "It is exciting that we have a path forward to identify new regeneration molecules."

More information: Evan L. Nichols et al, Functional Regeneration of the Sensory Root via Axonal Invasion, *Cell Reports* (2020). [DOI: 10.1016/j.celrep.2019.12.008](https://doi.org/10.1016/j.celrep.2019.12.008)

Provided by University of Notre Dame

Citation: New study identifies potential path forward for brachial plexus injury recovery (2020, January 16) retrieved 21 March 2023 from <https://medicalxpress.com/news/2020-01-potential-path-brachial-plexus-injury.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--