

# Wearable trackers estimate fitness levels without high-intensity exercise

February 14 2020

---



Credit: CC0 Public Domain

Researchers have developed a method to estimate cardiorespiratory fitness levels that could be applied to data captured by wearable fitness trackers during activities of daily life. This could facilitate testing for

those with low exercise tolerance and may reduce the need for medically supervised fitness testing. The study is published ahead of print in the *Journal of Applied Physiology*.

Cardiorespiratory fitness is the ability of the cardiovascular and respiratory systems to deliver enough oxygen to the muscles during physical activity. People with low [cardiorespiratory fitness](#) may have an increased risk of heart disease, heart attack or stroke. Testing the amount of oxygen the body uses during exercise (VO<sub>2</sub>max) can assess these risks and may also function as a [preventive measure](#). However, people must typically exercise to exhaustion to measure VO<sub>2</sub>max, and such testing requires medical supervision and specialized equipment. In addition, it may not be safe for all who need to undergo cardiorespiratory [fitness testing](#) to exercise at maximum effort. Methods that use lower intensity exercise (submaximal) do not always provide results as accurate as maximal testing.

"Assessment of [cardiorespiratory fitness] during everyday activities may be very promising to enable constant tracking of [cardiorespiratory fitness] in an unobtrusive manner, as it would simplify self-assessment by removing the need of exercise equipment typical of submaximal testing," an international team of researchers wrote.

The research team studied healthy adult volunteers between the ages of 18 and 55. The volunteers participated in a cycling test to measure their VO<sub>2</sub>max prior to the study. The researchers measured the volunteers' height, weight, fat and muscle mass and fitted them with a chest strap heart monitor and an activity monitor worn around the wrist. The volunteers were directed to wear the devices continuously during daytime hours for five consecutive days as they followed their typical daily routines.

Using a fitness index to calculate the relationship between energy

expenditure and [heart rate](#) during physical activity, the researchers predicted VO<sub>2</sub>max for each of the volunteers. Comparing the estimated VO<sub>2</sub>max with the true values from the cycling test, the margin of error was approximately 10 percent. This figure is lower than the mean error rate reported in previous research using submaximal methods to measure cardiorespiratory fitness. "This gives us confidence that the presented method offers valuable predictions of [cardiorespiratory fitness] useful for categorizing individuals of different fitness levels," the researchers wrote. "Future work should focus on extending the validity of the presented method with data to groups of elderly and patients."

"Cardio-respiratory [fitness](#) estimation from heart rate and body movement in daily life" is published ahead of print in the *Journal of Applied Physiology*.

**More information:** Alberto G. Bonomi et al. Cardio-respiratory fitness estimation from heart rate and body movement in daily life, *Journal of Applied Physiology* (2020). [DOI: 10.1152/jappphysiol.00631.2019](#)

Provided by American Physiological Society

Citation: Wearable trackers estimate fitness levels without high-intensity exercise (2020, February 14) retrieved 2 March 2023 from <https://medicalxpress.com/news/2020-02-wearable-trackers-high-intensity.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--