

Opioids for chronic non-cancer pain doubled in quarter century

25 February 2020



Credit: CC0 Public Domain

The number of people with chronic non-cancer pain prescribed an opioid medicine worldwide increased in the last two-and-a-half decades. But there was only a small number of studies reporting prescription data outside the United States, finds research led by the University of Sydney.

Chronic [pain](#) unrelated to cancer includes conditions such as chronic lower back pain, osteoarthritis and rheumatoid arthritis.

The researchers point to guidelines such as those from the Centres of Disease Control and Prevention in the United States that discourage the use of opioids to manage chronic non-[cancer pain](#) because of concerns about harmful effects and the lack of evidence about effectiveness. The use of opioids in the US and deaths from overdoses and addiction has been said to have reached epidemic proportions.

The systematic review of studies from across the world is the first to examine the literature about the extent opioid pain relievers are being prescribed to manage people with [chronic pain](#) conditions.

The findings are published today in the high-impact *Journal of Internal Medicine*.

The research spanned eight countries and evaluated 42 published studies that included 5,059,098 people with chronic pain conditions (other than cancer).

Two-thirds of the studies were from the US; one study was from Australia and the other studies were from the United Kingdom, Norway, India, Spain, Denmark and Canada.

Lead author Dr. Stephanie Mathieson from the University of Sydney's Institute for Musculoskeletal Health says that in the period 1991-2015, prescribing of opioid medicines increased markedly.

In the [early studies](#), opioid medicines were prescribed to about 20 percent of patients experiencing chronic pain but the later studies report rates of more than 40 percent.

"Over this period, on average around 30 percent of people with chronic pain were prescribed an opioid medicine," said Dr. Mathieson, from the School of Public Health in the Faculty of Medicine and Health.

"We noted that a higher proportion of people were prescribed a strong opioid medicine such as oxycodone compared to weak opioid pain-relieving medicines."

The authors also discovered there was insufficient data on the dose and duration of opioids prescribed to patients with chronic non-cancer pain.

Key findings:

Opioid prescribing:

Between 1991 and 2015, the researchers found in people with chronic pain (unrelated to cancer):

- Opioid prescribing increased over time from approximately 20 percent in early years to around 40 percent in later years.
- On average over this period approximately one in three patients (30.7 percent) were prescribed an opioid medicine.
- 42 percent of patients with chronic lower back pain were prescribed an opioid.
- The average age of those prescribed an opioid medicine was 55.7 years.
- Prescribing was not associated with the geographical location or the clinical setting where the opioids were prescribed (such as GPs or medical specialists).

University of Sydney; the University of Warwick, UK; the University of Notre Dame; the University of New South Wales and Monash University.

More information: *Journal of Internal Medicine* (2020). [DOI: 10.1111/joim.13026](https://doi.org/10.1111/joim.13026)

Provided by University of Sydney

Types of opioid painkillers:

In 17 studies that described the type of opioid pain relievers prescribed:

- 24.1 percent were strong combination products containing opioids (eg oxycodone plus paracetamol).
- 18.4 percent were strong opioids (eg oxycodone, morphine, fentanyl).
- 8.5 percent were weak opioids (eg codeine, tramadol).
- 11 percent were weak combination products containing opioids (eg codeine plus paracetamol).

An 'evidence' gap in global prescription data

The study aimed to establish a baseline for how commonly opioids are prescribed for people with chronic pain conditions (other than cancer). But the authors discovered a crucial evidence gap in [prescription data](#) in countries outside of the US.

"While we have sufficient data for this purpose for the US, we have little or no data for other countries," the authors write.

Dr. Mathieson says that studies in other countries, particularly low and [middle-income countries](#), are needed in order to check whether these countries are at risk of the problems seen in the US, where there is liberal use of [opioid](#) medicines.

This research is a collaboration between the

APA citation: Opioids for chronic non-cancer pain doubled in quarter century (2020, February 25)
retrieved 22 May 2021 from <https://medicalxpress.com/news/2020-02-opioids-chronic-non-cancer-pain-quarter.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.