

Breastfeeding linked to lower ovarian cancer risk

April 3 2020



Credit: iStock

An international study involving researchers from QIMR Berghofer Medical Research Institute has found women who breastfeed their



babies may lower their risk of developing ovarian cancer by almost 25 percent.

The research also shows the longer a woman breastfeeds, the greater the reduction in risk.

Senior Australian author and head of QIMR Berghofer's Gynaecological Cancers Group, Professor Penelope Webb, said breastfeeding was associated with a lower risk of developing all <u>ovarian cancers</u>, including the most lethal type called high-grade serous tumors.

"Overall, the risk of developing ovarian <u>cancer</u> dropped by 24 percent for <u>women</u> who breastfed, and even those who breastfed their children for three months or less had about an 18 percent lower risk of developing ovarian cancer," Professor Webb said.

"Mothers who breastfed their children for more than 12 months each had a 34 percent lower risk.

"Importantly, this benefit of breastfeeding lasted for at least 30 years after a woman stopped breastfeeding."

More than a thousand women died from ovarian cancer in Australia in 2019, accounting for almost five percent of female cancer deaths last year, according to the Australian Institute of Health and Welfare.

Only about 45 percent of women diagnosed with ovarian cancer survive at least five years after their diagnosis.

Professor Webb said the international study involved researchers from the Ovarian Cancer Association Consortium, who examined data from 9973 women with ovarian cancer and 13,843 control women from studies conducted around the world.



"Some past studies have linked breastfeeding to a reduced risk of ovarian cancer, but others found no association, so we wanted to look at this in a much bigger study to clarify the relationship," she said.

"The study results show a link between breastfeeding and reduced ovarian cancer rates, and reinforce the World Health Organization's recommendations that mothers should exclusively breast feed for at least six months if they can and continue doing so, with the addition of complementary foods, for two or more years.

"The research also shows that breastfeeding for even a short period of time may help reduce cancer risk.

"This study builds on previous work conducted at the Institute that found that breastfeeding was also associated with a <u>lower risk</u> of endometrial cancer (cancer of the womb)."

Professor Webb said more research is now needed to identify how <u>breastfeeding</u> affects <u>cancer risk</u>.

The study results have been published in the journal JAMA Oncology.

More information: Ana Babic et al. Association Between Breastfeeding and Ovarian Cancer Risk, *JAMA Oncology* (2020). <u>DOI:</u> <u>10.1001/jamaoncol.2020.0421</u>

Provided by QIMR Berghofer Medical Research Institute

Citation: Breastfeeding linked to lower ovarian cancer risk (2020, April 3) retrieved 25 February 2023 from <u>https://medicalxpress.com/news/2020-04-breastfeeding-linked-ovarian-cancer.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.