

## Too sick to work? Global survey captures behavior of healthcare and non-healthcare workers

13 May 2020



Credit: CC0 Public Domain

International survey shows that most workers would continue to work when sick with flu-like symptoms, according to a study published May 13, 2020 in the open-access journal *PLOS ONE* by Ermira Tartari from the International Society of Antimicrobial Chemotherapy Infection and Prevention Control (ISAC-IPC) Working Group and colleagues.

Working while sick is a reality for workers across sectors and countries, for reasons including lack of paid sick leave, work culture, and individual judgement. Against the backdrop of the current COVID-19 crisis, this behavior clearly warrants challenging—especially as it manifests in the healthcare industry, considering the potential for

transmission of respiratory disease between patients and <u>healthcare staff</u>.

In order to better understand the phenomenon of working while sick for <a href="healthcare workers">healthcare workers</a> and non-healthcare workers alike, Tartari and colleagues conducted an opt-in cross-sectional online survey of the members and international networks of the International Society of Antimicrobial Chemotherapy (ISAC) Infection Prevention and Control (IPC) Working Group from October 2018-January 2019.

In total, 533 respondents from 49 countries participated—of these, 249 were healthcare workers (46.7 percent) and 284 non-healthcare workers (53.2 percent). A clear majority from both groups (99.2 percent of healthcare workers and 96.5 percent of non-healthcare workers) would work through "minor" symptoms like a sore throat, sneezing/runny nose, or cough. 58.5 percent of respondents stated they'd continue to work when sick with an influenza-like illness (including major symptoms like muscle aches and fever), with no significant variation between healthcare workers and non-healthcare workers.

Healthcare workers did significantly diverge in some ways: 26.9 percent of healthcare workers would continue to work even when experiencing a fever (one of the most major symptoms), compared with 16.2 percent of non-healthcare workers; similarly, only 45.8 percent of healthcare workers would avoid a colleague exhibiting influenza-like symptoms, compared with 60.9 percent of non-healthcare workers. Healthcare workers were also more willing to receive the influenza vaccine (81.1 percent, versus 56.7 percent of non-healthcare workers).

Though this study is limited by the fact that the



participants were self-selected, with 90 percent of respondents coming from high-income countries, the results are concerning, since over half of both healthcare and non-healthcare workers report being willing to continue working through major influenza symptoms, despite international guidelines. In the midst of a pandemic, future strategies to prevent the transmission of influenza-like illnesses in work settings—especially healthcare, where workers often care for immunocompromised patients—are crucial, and the authors note that a broad cultural change is required, in addition to sufficient sick leave and access to the influenza vaccine.

The authors add: "At the time of the present COVID pandemic it is important to realize that healthcare workers, despite feeling moderately sick during flu seasons, feel the obligation to work."

**More information:** Tartari E, Saris K, Kenters N, Marimuthu K, Widmer A, Collignon P, et al. (2020) Not sick enough to worry? "Influenza-like" symptoms and work-related behavior among healthcare workers and other professionals: Results of a global survey. *PLoS ONE* 15(5): e0232168. doi.org/10.1371/journal.pone.0232168

Provided by Public Library of Science
APA citation: Too sick to work? Global survey captures behavior of healthcare and non-healthcare workers (2020, May 13) retrieved 16 September 2022 from <a href="https://medicalxpress.com/news/2020-05-sick-global-survey-captures-behavior.html">https://medicalxpress.com/news/2020-05-sick-global-survey-captures-behavior.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.