

# Infection risk up in month after CAR T-cell immunotherapy

29 May 2020



infections were bacteremias or respiratory viral infections (39 and 43 percent, respectively). Prior hematopoietic cell transplantation, immunoglobulin G (IgG) level

(HealthDay)—For children and young adults receiving CD19 chimeric antigen receptor (CAR) T-cell infusion (CTI), infection rates increase in the first month after treatment and then decrease, according to a study published in the May issue of *Open Forum Infectious Diseases*.

Surabhi B. Vora, M.D., M.P.H., from the University of Washington in Seattle, and colleagues reviewed the [medical records](#) of 83 patients 26 years receiving CTI at a single institution between 2014 and 2017. The number of infections per 100 days at risk was calculated in the 90 days preceding and 0 to 28 and 29 to 90 days after CTI.

The researchers found that 98 percent of patients had refractory or relapsed [acute lymphoblastic leukemia](#). In the 90 days before CTI, infections occurred in 54 percent of patients (infection density, 1.23) compared with 40 percent of patients in the 28 days following CTI (infection density, 2.89). In the 29 to 90 days after CTI, infection density decreased to 0.55. Most

APA citation: Infection risk up in month after CAR T-cell immunotherapy (2020, May 29) retrieved 9 September 2022 from <https://medicalxpress.com/news/2020-05-infection-month-car-t-cell-immunotherapy.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*