

# Physical activity of older people requires tailored monitoring

July 8 2020

---



Commercial activity monitors may underestimate the exertion level of older adults' activity. Credit: University of Jyväskylä

The ability to move about may deteriorate when aging, and must be considered when assessing physical activity in older people. A study on

active aging at the University of Jyväskylä examined movement that exceeds the intensity of preferred walking speed.

Improving physical performance requires exercising regularly beyond one's usual level of exertion. The body then adapts to the new level of exertion by improving performance. Many [activity](#) monitors on the market have been developed for young and middle-aged people who have higher physical performance than [older adults](#). Therefore, activity monitors may underestimate the exertion level of older adults' activity.

In the study at the University of Jyväskylä, preferred walking speed was measured in a six-minute walking test. In addition, the participants wore an activity monitor while living their day-to-day life.

"By measuring their preferred walking speed we were able to assess the time that our participants exercised more strenuously than what is their usual exertion level and what is beyond their [comfort zone](#)," explains postdoctoral researcher Laura Karavirta from the Gerontology Research Center and Faculty of Sport and Health Sciences.

The participants in the study accumulated 62 minutes of activity, on average, beyond the intensity of their preferred walking speed. Interestingly, the amount of activity was similar in 75-, 80- and 85-year-old people, regardless of age.

"The new method enables us to investigate [physical activity](#) as individual behavior, which is not influenced by fitness level," Karavirta explains. "A physically [active lifestyle](#) is about challenging oneself according to one's own abilities. Light intensity movement is also important, but at least moderate exertion is required for improving physical performance."

The prevailing recommendation for all adults is a minimum of 150

minutes of at least moderate intensity physical activity per week. The general definition for moderate intensity is equivalent to exceeding three times the energy consumption of rest. Individual exertion at this intensity varies according to person's fitness level.

"For most [young adults](#), it feels easy and corresponds to slow walking but for some older adults it may be the hardest effort they can perform," Karavirta says.

**More information:** Laura Karavirta et al. Individual Scaling of Accelerometry to Preferred Walking Speed in the Assessment of Physical Activity in Older Adults, *The Journals of Gerontology: Series A* (2020). [DOI: 10.1093/gerona/glaa142](https://doi.org/10.1093/gerona/glaa142)

Provided by University of Jyväskylä

Citation: Physical activity of older people requires tailored monitoring (2020, July 8) retrieved 10 April 2023 from <https://medicalxpress.com/news/2020-07-physical-older-people-requires-tailored.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------