

# Mom and baby share 'good bacteria' through breast milk

July 10 2020, by Kerry Blackadar

---



A new study has found that bacteria are shared and possibly transferred from a mother's milk to her infant's gut, and that breastfeeding directly at the breast best supports this process. Credit: Kevin Liang/Unsplash

A new study by researchers at the University of British Columbia and

the University of Manitoba has found that bacteria are shared and possibly transferred from a mother's milk to her infant's gut, and that breastfeeding directly at the breast best supports this process.

The research, published today in *Cell Host & Microbe*, found that certain bacteria, including *Streptococcus* and *Veillonella*, co-occur in mothers' [milk](#) and their infants' stool, and this co-occurrence is higher when infants nurse directly at the breast.

"Our study confirms that [breast milk](#) is a major driver of infant gut microbiota development," said the study's senior co-author Dr. Stuart Turvey, a professor in UBC's department of pediatrics and investigator at BC Children's Hospital. "We found that breastfeeding exclusivity and duration was strongly associated with a baby's overall gut microbiota composition and that breast milk bacteria shape a baby's [gut microbiome](#) to a similar degree as other known modifiers of the gut microbiota such as birth mode—meaning a cesarean-section or vaginal delivery."

According to the researchers, this is the first study to evaluate the association of multiple breast milk feeding practices (mode, exclusivity, and duration), milk bacteria, and milk components with infant gut microbiota composition at multiple time points in a baby's first year.

The researchers analyzed the microbiome of infants' stool and their mothers' breastmilk using 16S rRNA sequencing, a technique used to identify, classify and determine the abundance of microbes.

The 1,249 mother-baby pairs involved in the research are participating in the CHILD Cohort Study (CHILD), a world-leading birth cohort study in maternal, newborn and child health research. The findings build upon previous CHILD research that showed pumping breast milk is associated with differences in both milk microbiota composition and infant health.

"Uniquely, our study showed that while breast milk and the infant gut have distinct microbiota compositions, there are a few commonly shared bacteria that were more prevalent and abundant in breast milk of mothers who only nursed directly at the breast, while other bacteria showed dose-dependent associations with exclusive breastfeeding," said the study's senior co-author Brett Finlay, professor in the departments of biochemistry and [molecular biology](#), and microbiology and immunology at UBC.

"These results advance the hypothesis that [breast](#) milk may act as an incubator that enriches, protects and transports certain bacteria to a baby's [intestinal tract](#) and this may give us clues about which bacteria could make good probiotics since they appear to withstand the trip to the baby's gut," Finlay added.

**More information:** Kelsey Fehr et al, Breastmilk Feeding Practices Are Associated with the Co-Occurrence of Bacteria in Mothers' Milk and the Infant Gut: the CHILD Cohort Study, *Cell Host & Microbe* (2020). [DOI: 10.1016/j.chom.2020.06.009](https://doi.org/10.1016/j.chom.2020.06.009)

Provided by University of British Columbia

Citation: Mom and baby share 'good bacteria' through breast milk (2020, July 10) retrieved 5 July 2023 from <https://medicalxpress.com/news/2020-07-mom-baby-good-bacteria-breast.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--