


# Scientists observe learning processes online in the brain

July 21 2020

---



Credit: Pixabay/CC0 Public Domain

Stimulating the fingertip rhythmically for a sustained period of time markedly improves touch sensitivity. A research team led by Associate Professor Dr. Hubert Dinse at Ruhr-Universität Bochum (RUB) analyzed the impact of this process in the brain. Using electroencephalography (EEG), the scientists recorded neuronal activity

of brain areas associated with tactile processing. They were able to observe changes in activity over time—possibly illustrating a learning process. The team reported their findings in *Frontiers in Human Neuroscience*.

## **Learning through repetition**

In daily life, people learn through practice and repetition, enabled by a [brain](#) process called neuronal plasticity. A prominent example of a cellular basis for such plasticity processes is called long-term potentiation, the ability of neurons to increase communication efficiency with other neurons they are connected to.

In analogy to the rhythmic neuronal activity underlying long-term potentiation, the RUB team has developed a method of stimulation-based learning. Thereby senses, such as the sense of sight or touch, are rhythmically stimulated. A well-studied example of this is electrical stimulation of the fingertips, which—if administered at the correct frequency—has been shown to increase tactile sensitivity in the stimulated fingertip. Studies show that this fingertip stimulation leads to significant plasticity processes in the somatosensory cortex. However, it has not yet been proven whether long-term potentiation is the foundation of these processes.

## **Two groups, two experiments**

The RUB neuroscientists have studied stimulation-based learning in volunteers with EEG recordings. Their goal was to assess neuronal activity as well as its development during this learning process. As such, they carried out two experiments with two groups of subjects. The first experiment served as a control study to confirm that the method of air stimulation administered through an inflatable membrane used in this

experiment has the same positive effect on touch sensitivity as the established method of electrical stimulation. The electrical version could not be used in this case, as it distorts the signal of EEG recordings with electrical artifacts.

In the second experiment, volunteers received the aforementioned painless air-stimulation on their fingertip for 40 minutes. At the same time, the activity in the somatosensory cortex of the test subjects was measured with EEG. The scientists concentrated on the area of the brain associated with sensory processing of the hand.

## **Nerve cells adapt their activity**

"Using electroencephalic measurements of brain activity, we were able to show that large cell ensembles adapt their activity to the frequency of the stimulation during the active stimulation phases. This reaction remains stable over 20 minutes, without any signs of habituation, which is very similar to cellular long-term potentiation," explains Dr. Marion Brickwedde, first author of the study.

Furthermore, the scientists were also able to observe how neuronal responses to the stimulation changed over time. They found that the shape of event-related potentials, which represent stimulus processing in the brain, was changing. The event-related desynchronization of the alpha rhythm, a typical response to tactile stimuli, was also reduced after 20 minutes.

"These processes may represent excitability changes in tactile brain areas, that is, a directly observable [learning process](#). It is not yet possible to draw the definite conclusion that the here-applied finger stimulation actually triggers long-term potentiation in the human sensorimotor cortex. But taking into account previous findings which show, for example, the dependence of both processes on the same neuronal

receptor type, the accumulating evidence speaks volumes," explains Marion Brickwedde.

**More information:** Marion Brickwedde et al. 20 Hz Steady-State Response in Somatosensory Cortex During Induction of Tactile Perceptual Learning Through LTP-Like Sensory Stimulation, *Frontiers in Human Neuroscience* (2020). [DOI: 10.3389/fnhum.2020.00257](https://doi.org/10.3389/fnhum.2020.00257)

Provided by Ruhr-Universitaet-Bochum

Citation: Scientists observe learning processes online in the brain (2020, July 21) retrieved 13 February 2024 from <https://medicalxpress.com/news/2020-07-scientists-online-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.