


Calcium channel subunits play a major role in autistic disorders

July 22 2020



After 21 days in vitro, hippocampal neurons of the rat form many synaptic contacts and communicate with each other. Within the study such neuronal networks cultivated on multi-electrode arrays were used to invest the

synaptogenic function of $\alpha 2\delta$ -subunits. Credit: ill. by Artur Bikbaev

The ability of the human brain to process and store information is determined to a large extent by the connectivity between nerve cells. Chemical synapses are very important in this context as they constitute the interface for the transmission of information between individual nerve cells. Abnormalities in the formation of synapses cause many neurological disorders such as autism. Neurobiologists at Johannes Gutenberg University Mainz (JGU) have found new evidence that specific calcium channel subunits play a crucial role in the development of excitatory and inhibitory synapses.

$\alpha 2\delta$ subunits have different effects on the formation of new synapses

Autism spectrum disorder is a condition involving impaired development that begins with birth and is usually manifested when the individual in question exhibits difficulties in social interaction and communication. It is postulated that the main underlying cause is disruption of synapse-mediated interaction between nerve cells.

The results of several studies indicate that so-called $\alpha 2\delta$ subunits of calcium channels are involved in the formation and fine-tuning of excitatory and inhibitory nerve cells, but little has been known to date about when and how specifically the four forms of $\alpha 2\delta$ subunits are involved. It is this aspect that the research team led by Professor Martin Heine of the Institute of Developmental Biology and Neurobiology at Mainz University has now addressed. What is particularly interesting is their research finding that the two dominant $\alpha 2\delta$ subunits in the hippocampus, $\alpha 2\delta 1$ and $\alpha 2\delta 3$, have different effects on synaptogenesis in [neuronal networks](#).

In order to investigate the underlying mechanism, the researchers prepared isolated networks of hippocampal neurons. The results show that during the early phase of the development of neural networks, subunit $\alpha 2\delta 3$ promotes the release of an inhibitory neurotransmitter, triggers the formation of [inhibitory synapses](#), and boosts the growth of axons from inhibitory neurons. "The $\alpha 2\delta 3$ subunit is obviously an important factor with regard to the early development of neural networks," explained Heine. At later development phases and in more mature neuronal networks, it is subunit $\alpha 2\delta 1$ that fosters excitatory stimulus transmission and synaptogenesis.

Connectivity relies on concerted cooperation between $\alpha 2\delta 1$ and $\alpha 2\delta 3$

In their article in *The Journal of Neuroscience*, the researchers proposed "that formation of connectivity in neuronal networks is associated with a concerted interplay of $\alpha 2\delta 1$ and $\alpha 2\delta 3$ subunits of calcium channels". Dr. Artur Bikbaev, one of the lead authors from JGU, further concluded that the calcium channel subunits are molecules that are relevant to the development of the brain. New data has confirmed the assumption that there is a link between an aberration in the genes that code for subunits $\alpha 2\delta 1$ and $\alpha 2\delta 3$ and autistic spectrum disorders. An imbalance in the ratio of excitatory to inhibitory neurons is also thought to be the cause of the epileptic seizures which very frequently accompany autism spectrum disorder.

In addition to the team at JGU, researchers at the Leibniz Institute for Neurobiology, the University of Münster, and the Medical University of Innsbruck were also involved in the project.

More information: Arthur Bikbaev et al, Auxiliary $\alpha 2\delta 1$ and $\alpha 2\delta 3$ Subunits of Calcium Channels Drive Excitatory and Inhibitory Neuronal Network Development, *The Journal of Neuroscience* (2020). [DOI:](#)

[10.1523/JNEUROSCI.1707-19.2020](https://doi.org/10.1523/JNEUROSCI.1707-19.2020)

Provided by Universitaet Mainz

Citation: Calcium channel subunits play a major role in autistic disorders (2020, July 22)
retrieved 28 December 2022 from

<https://medicalxpress.com/news/2020-07-calcium-channel-subunits-major-role.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.