

Hearing loss linked to neurocognitive deficits in childhood cancer survivors

July 30 2020



Research shows that severe hearing loss in childhood cancer survivors is associated with neurocognitive deficits independent of type of therapy. Credit: St. Jude Children's Research Hospital

Scientists at St. Jude Children's Research Hospital are studying how

hearing loss can affect the neurocognitive abilities of childhood cancer survivors. Findings show that survivors with severe hearing loss are at a significant increased risk for neurocognitive deficits, independent of what type of therapy they receive. The work appears as an advance online publication today in *JAMA Oncology*.

"There has been a lot of research on how children who are born deaf can be affected academically and psychosocially, but far less for kids who acquire [hearing](#) loss at some point during their childhood," said corresponding author Johnnie Bass, Ph.D., of St. Jude Rehabilitation Services. "Our goal was to report on the prevalence, severity and risk of hearing loss in a large cohort of cancer survivors to assess the impact of hearing impairment on neurocognitive function."

The researchers evaluated 1,520 childhood cancer survivors and found that more than one-third had severe hearing loss. Those survivors with severe hearing loss were found to be at increased risk for neurocognitive deficits. This effect was independent of having been exposed to neurotoxic therapy, when compared to survivors with normal hearing or mild hearing loss.

This study is the first to objectively measure hearing and neurocognitive function in a large cohort of long-term survivors of childhood cancer stratified by treatment exposures.

"Even patients not exposed to neurotoxic therapies who develop mild hearing deficits can have problems with their neurocognitive skills," said senior author Kevin Krull, Ph.D., of St. Jude Epidemiology and Cancer Control. "This makes it important to identify these patients early and suggest interventions to help improve their hearing and thus their neurocognitive outcomes."

Hearing aids are often recommended to assist with treatment-related

hearing loss. The researchers found that of 330 survivors with severe hearing loss in the study for whom a hearing aid had been previously recommended, only 23% were consistently using the devices. Research in individuals who are born with hearing loss, as well as in the elderly with hearing loss, suggests that hearing aids can improve neurocognitive outcomes. More research is needed to determine if this is the case for childhood cancer survivors and to better understand barriers that prevent hearing aid use.

The work relied on hearing assessments gathered through the St. Jude Lifetime Cohort study (St. Jude LIFE). St. Jude LIFE brings long-term childhood cancer survivors back to the hospital for regular health screenings throughout their adult lives. Findings from the study help [childhood cancer survivors](#) learn more about their health needs, while providing novel insights into the late effects of [childhood cancer](#) treatment.

More information: *JAMA Oncology* (2020).

jamanetwork.com/journals/jamao.../jamaoncol.2020.2822

Provided by St. Jude Children's Research Hospital

Citation: Hearing loss linked to neurocognitive deficits in childhood cancer survivors (2020, July 30) retrieved 4 February 2024 from

<https://medicalxpress.com/news/2020-07-loss-linked-neurocognitive-deficits-childhood.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--