


Cardiovascular risk factors tied to COVID-19 complications and death

August 14 2020



Cardiovascular risk profile of hospitalized COVID-19 patients. Each line represents the result of the meta-analysis for a single endpoint. The square represents the summary effect size (proportion) and the horizontal line the relative 95% Confidence Interval. Credit: Sabatino et al, 2020 (*PLOS ONE*, CC BY 4.0)

COVID-19 patients with cardiovascular comorbidities or risk factors are more likely to develop cardiovascular complications while hospitalized, and more likely to die from COVID-19 infection, according to a new

study published August 14, 2020 in the open-access journal *PLOS ONE* by Jolanda Sabatino of Universita degli Studi Magna Graecia di Catanzaro, Italy, and colleagues.

For most people, the Novel Coronavirus Disease 2019 (COVID-19) causes mild illness, however it can generate severe pneumonia and lead to death in others. It is crucial for clinicians working with cardiovascular patients to understand the clinical presentation and [risk factors](#) for COVID-19 infection in this group.

In the new study, researchers analyzed data from 21 published [observational studies](#) on a total of 77,317 hospitalized COVID-19 patients in Asia, Europe and the United States. At the time they were admitted to the hospital, 12.89% (95% CI 8.24-18.32) of the patients had cardiovascular comorbidities, 36.08% (95% CI 20.25-53.64) had hypertension and 19.45% (95% CI 12.55-27.45) had diabetes.

Cardiovascular complications were documented during the [hospital stay](#) of 14.09% (95% CI 10.26-20.23) of the COVID-19 patients. The most common of these complications were arrhythmias or palpitations; significant numbers of patients also had myocardial injury. When the researchers analyzed the data, they found that pre-existing cardiovascular comorbidities or risk factors were significant predictors of [cardiovascular complications](#) ($p=0.019$), but age ($p=0.197$) and gender ($p=0.173$) were not. Both age and pre-existing cardiovascular comorbidities or risk factors were significant predictors of death.

The authors add: "Cardiovascular complications are frequent among COVID-19 patients and might contribute to adverse clinical events and mortality."

More information: Sabatino J, De Rosa S, Di Salvo G, Indolfi C (2020) Impact of cardiovascular risk profile on COVID-19 outcome. A

meta-analysis. *PLoS ONE* 15(8): e0237131.
doi.org/10.1371/journal.pone.0237131

Provided by Public Library of Science

Citation: Cardiovascular risk factors tied to COVID-19 complications and death (2020, August 14) retrieved 3 April 2023 from
<https://medicalxpress.com/news/2020-08-cardiovascular-factors-tied-covid-complications.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.