

Study suggests opioid use linked to pregnancy loss, lower chance of conception

18 August 2020



Credit: CC0 Public Domain

Opioid use among women trying to conceive may be associated with a lower chance of pregnancy, suggests a National Institutes of Health study. Moreover, opioid use in early pregnancy may be associated with a greater chance of pregnancy loss. The study appears in *Epidemiology*.

"Our findings indicate that [women](#) who are pregnant or planning a [pregnancy](#) should, along with their physicians, consider the potential effects opioids may have on their ability to conceive or sustain a pregnancy," said Kerry Flannagan, Ph.D., the primary author of the study and a postdoctoral researcher in the Division of Intramural Population Health Research at NIH's Eunice Kennedy Shriver National Institute of Child Health and Human Development.

According to the authors, much of the research on prescription [opioid use](#) has focused on the effects of drug dependency. Little information exists on non-habitual, periodic [opioid](#) use around the time of conception and early in pregnancy.

The researchers analyzed data from the Effects of

Aspirin in Gestation and Reproduction (EAGeR) trial, which investigated low-dose aspirin as a treatment to prevent pregnancy loss. Participants were women from 18 to 40 years old with a history of one or two pregnancy losses. Women were followed for six monthly cycles if they did not get pregnant and throughout pregnancy if they did. The women provided [urine samples](#), which were analyzed for various prescription opioids.

Of the 1,228 women in the study, 226 (18%) had used opioids while trying to conceive and 33 (5%) of 685 women who became pregnant had used opioids in early pregnancy. None tested positive for methadone or buprenorphine, typically used to treat opioid dependence.

Opioid use before conception was associated with a 29% lower chance of achieving pregnancy during a given monthly cycle, compared to women who had not used opioids. Among the women who became pregnant, those who used opioids around the time of conception were 1.5 times as likely to have a miscarriage as women who had not. Women who used opioids in the first four weeks of pregnancy were more than twice as likely to have a miscarriage. Women who used opioids in weeks four through eight of pregnancy were 2.5 times as likely to have a miscarriage.

The authors called for additional research on how opioid use affects fertility and early pregnancy. They added that until more is known, patients and physicians should evaluate the potential risks and benefits of opioids for [pain management](#) among women who are pregnant or may become pregnant, including those undergoing assisted reproduction procedures that may involve opioid treatment to manage pain.

More information: Kerry S. Flannagan et al. Is opioid use safe in women trying to conceive?, *Epidemiology* (2020). DOI: [10.1097/EDE.0000000000001247](https://doi.org/10.1097/EDE.0000000000001247)

Provided by NIH/Eunice Kennedy Shriver National
Institute of Child Health and Human Development

APA citation: Study suggests opioid use linked to pregnancy loss, lower chance of conception (2020, August 18) retrieved 19 October 2022 from <https://medicalxpress.com/news/2020-08-opioid-linked-pregnancy-loss-chance.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.