

Study tackles long-term opioid use among retired NFL players

16 September 2020



The investigators found that nearly 50% of the former players who reported using opioids in 2010 were still using them in 2019. About 60% of those who used opioids in 2010 reported moderate to severe depressive symptoms in the past week, and had lower scores on a measure of mental health-

related quality of life.

continue using them.

(HealthDay)—Many former National Football League (NFL) players who took opioid painkillers early in their retirement still used them nine years later, a new study finds.

The researchers also found that those who continued to use opioids were more likely to report moderate to severe depressive symptoms and low mental health-related quality of life.

Long-term opioid use among former NFL players may be due to a number of factors, said study coauthor Zachary Mannes, a recent graduate of the doctoral program in clinical psychology at the University of Florida.

"It's hard to imagine that bodily pain doesn't contribute to continued opioid use, as retired NFL athletes may rely on opioids in an effort to manage chronic pain that has developed as a result of an accumulation of orthopedic injuries during an NFL career," Mannes said in a university news release.

Opioids can be effective for acute pain control, but their use isn't recommended for chronic pain. Longterm use can lead to greater pain sensitivity and reduce the likelihood of chronic pain recovery, explained Mannes, now a postdoctoral fellow at the Columbia University Mailman School of Public Health in New York City.

Previous research has shown that NFL players are often overprescribed opioids to control pain and that many retired professional football players

In this study, the researchers interviewed 90 former NFL athletes who'd taken part in a 2010 study examining the prevalence and risk factors of opioid

use and misuse among NFL retirees.

The former NFL players might also be using opioids as a way to cope with mental health issues.

"It is possible that NFL retirees with depressive symptoms are more likely to self-medicate their mental health symptoms and emotional pain through use of opioids," Mannes said.

He added that <u>medical professionals</u> working with retired NFL athletes should monitor <u>opioid use</u> and consider alternative pain management strategies, such as exercise, physical rehabilitation and psychotherapy.

The study will be published in the October issue of the journal *Drug and Alcohol Dependence*.

More information: The U.S. Food and Drug



Administration has more on opioid medications.

Copyright © 2020 HealthDay. All rights reserved.

APA citation: Study tackles long-term opioid use among retired NFL players (2020, September 16) retrieved 16 July 2022 from https://medicalxpress.com/news/2020-09-tackles-long-term-opioid-nfl-players.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.