

# Videos most effective in communicating with parents about secondhand smoke risks

30 September 2020



Credit: Unsplash/CC0 Public Domain

The best way to communicate with parent smokers about the risks of secondhand smoke to their children is to use videos depicting the risks, as well as solutions to reduce those risks.

Those were the findings of a study, published in the *Journal of Health Communication*, which demonstrates that video messages, compared to text-only messages, are significantly more effective in influencing parent smokers' intentions to protect their offspring from exposure to [secondhand smoke](#).

The effects and diseases caused by secondhand smoking kills 41,000 people in America annually (Centers for Disease Control and Prevention, 2017). Of the 58 million Americans exposed to secondhand smoke, infants and [children](#) are the most vulnerable to adverse [health](#) effects.

In order to assess which format of communication may be most effective to getting people to not only stop smoking, but stop smoking near their children, experts from both the Center of Tobacco Studies at Rutgers University and The SIDS Center of New

Jersey, assessed 623 adult daily smokers living with children in the US, aged 0–15.

The participants were assigned to view messages that differed by recommendation (stopping altogether—known as cessation vs cessation plus exposure reduction) and format (video vs text-only), or to a no-message control group.

All messages delivered to participants discussed the health risks of secondhand smoke. The cessation messages encouraged smokers to seek information about getting help to quit and for couples to work together to help each other quit, while the "cessation plus exposure reduction" messages additionally discussed strategies for parent smokers to protect their children from secondhand smoke, like smoking outdoors, washing toys, surfaces, and clothes exposed to secondhand smoke, and not smoking in a car with children.

Results showed:

- Parent smokers who saw video messages had significantly greater intentions to reduce secondhand smoke exposure to their children, as well as discuss the issue and risks of secondhand smoke with other parents, compared to parent smokers that saw text-only messages.
- Parents who saw videos also had significantly greater intentions to quit smoking altogether and get help to quit smoking compared to parents who did not view a message.
- Parent smokers who viewed either message recommendation reported greater harm perceptions, perceptions of self-efficacy to protect children from secondhand smoke, and intentions to get help to quit smoking than the no-message control group.
- Cessation plus exposure reduction

recommendations elicited greater quit intentions than the no-message control.

Reporting on their findings, lead author Dr. Jennah Sontag stated, "All forms of communication tested in this study were effective in influencing perceptions and intentions about the risks of secondhand smoke for children, indicating that even communication that lasts only one to two minutes or is presented in text form may influence parent smokers to reduce or eliminate secondhand smoke exposure to children.

"But our research suggests that sharing this information visually through a video can be especially effective, and these videos should include recommendations for both cessation and ways to protect children from secondhand smoke.

"We already know that visual messages are better than text-only formats of tobacco-related risk communication. However, this study clearly demonstrates that the visual portrayals of the potential health risks of secondhand smoke exposure among infants and children, from asthma, ear infections, respiratory problems or SIDS, may have influenced greater harm perceptions due to the severity of these outcomes."

It is estimated that two in five children in the US are exposed to secondhand smoking. However, notable disparities exist including those from low-income communities, 3- to 11-year olds, and black children, who are at highest risk of exposure. Several environmental factors play a role in this exposure, including a lack of smokefree laws in public spaces or smokefree rules at home, and living in multiunit housing. US healthcare expenditures of secondhand smoke exposure among children living in public housing were more than \$180 million in 2011 alone.

Only 55% of smoking mothers quit smoking during pregnancy, and 70% of these relapse into smoking after childbirth.

Sontag believes these types of videos are more likely to reach wider audiences—due to their shareability—and therefore should be considered for future anti-smoking and secondhand smoking

communications.

"Video shows the greatest potential to influence perceptions and intentions related to reducing secondhand smoke exposure. To reach parent smokers, such messaging can be used by practitioners in clinical settings, such as waiting rooms and examination rooms, which allow for follow-up discussions, or in family-based education and community organizations. Because videos are more appropriate for the digital/social media landscape, messages can be further disseminated among [parents](#) and other caregivers," she adds.

The paper calls for future research to assess the extent to which secondhand [smoking](#)-related communication that uses videos and includes recommendations for cessation and exposure reduction influences parent smokers to inquire with their doctor about cessation options. "It is also important to assess whether strategies that protect children from secondhand smoke are carried out long term by parent smokers who do not quit," it states.

Limitations of the study include that over 70% of participants were female. In addition, parent smokers viewed only one message one time on a computer screen in a controlled experimental setting. Messages viewed in a natural environment (e.g., clinics, hospitals) and multiple exposures may produce different results. Additionally, 38% of participants reported that they never smoke inside the home, implying that these participants are already exhibiting one of the exposure-reduction behaviors that reduces secondhand smoke exposure to children; recruitment of only parent smokers that always smoke inside the home may have produced different results.

**More information:** Jennah M. Sontag et al, Secondhand Smoke Risk Communication: Effects on Parent Smokers' Perceptions and Intentions, *Journal of Health Communication* (2020). [DOI: 10.1080/10810730.2020.1797947](https://doi.org/10.1080/10810730.2020.1797947)

Provided by Taylor & Francis

APA citation: Videos most effective in communicating with parents about secondhand smoke risks (2020, September 30) retrieved 11 November 2022 from <https://medicalxpress.com/news/2020-09-videos-effective-parents-secondhand.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*