


Remembering novelty: Researchers work to better understand process of memory

October 15 2020



Layers of the hippocampus. The mossy cell bodies are red, interneuron cells connecting neurons are green, and other neurons are blue. Credit: Ryuichi Shigemoto

Imagine going to a café you have never been to. You will remember this



new environment, but when you visit it again and again fewer new memories about the environment will be formed, only the things that changed will be really memorable. How this long-term memory are regulated is still not fully understood. Ryuichi Shigemoto from the Institute of Science and Technology Austria (IST Austria) in cooperation with researchers from Aarhus University and the National Institute for Physiological Sciences in Japan now have uncovered a new keystone in the formation of memories. In their study published in *Current Biology*, they investigated a signaling path in the hippocampus area of the brain and showed how it controls making new memories about experiencing new environments.

The hippocampus is a central area in the brain that plays an important role in transferring information from the short-term memory to the longterm memory. Of the many interlocking parts of the hippocampus, the researchers focused on the connection between the so-called mossy cells that receive novelty signals of sensory input about the environment and the so-called granule cells to which this information is relayed. In diseases like Alzheimer's this part of the brain is one of the first ones affected.

Watching Neurons

The scientists used four different approaches for this study in order to rigorously investigate these new findings. First, they put the hippocampus under the microscope and studied the structure of how the mossy cells are connected to the granule cells showing their many complex connections.

Second, they used the technique of calcium imaging that allows live monitoring of neuron activity as these genetically modified cells light up when activated. When exposing the <u>animals</u> a new environment for several days, the activity of the mossy cells sending signals to the granule



cells first was high and then became less and less. When they then put the mice into another new environment, the activity sprung up again, therefore showing that these neurons are specifically relevant to process new environmental input.

Third, the researchers followed traces in the neuron left by the signals. Neural activity in these cells triggers the expression of a certain gene, meaning the production of the corresponding protein that is encoded in it. The more activity there was, the more of this protein they can find afterwards. In the granule cells they found production of this protein, which correlated with activity of the mossy cells.

Dreading New Places

And lastly, the scientists used behavioral studies to see the effects of this pathway in the hippocampus on memory formation. This is especially important, because the connection between memory formation and behavior can tell them a lot about the brain's functions. They combined a negative sensory stimulus, a small electric shock, with putting the animals in a new environment. The mice then quickly learned to associate the new environment with unpleasant feelings and their negative reaction of freezing on the spot was measured.

When the researches used drugs to inhibit the activity of the mossy cells—the ones receiving the signals about the new environment—and then did the negative conditioning, the mice did not remember the connection between the new environment and the unpleasant feeling. Additionally, when the animals were first accustomed to the new environment and then conditioned, there was also no activation of the mossy cells, and therefore no association between the environment and the shocks.

On the other hand, if the mossy cells were artificially activated, this



association could be formed even after the animals were already used to the new environment. This clearly shows how the mossy <u>cells</u> in the hippocampus react to novel input and trigger the formation of new longterm memories in mice.

Next Steps in Understanding

Whether the exact same processes happen in the human brain is still an open question, but these new findings are an important first step in understanding this part of our most complex organ. Ryuichi Shigemoto and his collaborators are conducting <u>fundamental research</u> that may one day help to address degenerative brain diseases that affect memory formation, but this is still a while away.

He cautiously states: "This <u>research field</u> is very competitive and new findings arise quickly. There are many disputed mechanisms on <u>memory</u> formation, but our findings corroborate an existing hypothesis and are very robust, thus opening up a new field of neuroscience research and furthering our understanding of the brain."

Animal welfare

Understanding how the <u>brain</u> stores and processes information is only possible by studying the brains of animals while they carry out specific behaviors. No other methods, such as in vitro or in silico models, can serve as alternatives. The animals were raised, kept and treated according to the strict regulations of Austrian law.

More information: Felipe Fredes et al. Ventro-dorsal Hippocampal Pathway Gates Novelty-Induced Contextual Memory Formation. *Current Biology*. Published:October 15, 2020. <u>DOI: 10.1016/j.cub.2020.09.074</u>



Provided by Institute of Science and Technology Austria

Citation: Remembering novelty: Researchers work to better understand process of memory (2020, October 15) retrieved 5 May 2023 from <u>https://medicalxpress.com/news/2020-10-novelty-memory.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.