

# 'Vanished' or 'hidden' prostate cancer? Men with negative biopsies during active surveillance have good outcomes

17 November 2020

Can early-stage prostate cancer "vanish" during follow-up? More likely the cancer is just "hidden"—either way, negative biopsies during active surveillance for prostate cancer are associated with excellent long-term outcomes, reports a study in *The Journal of Urology*, an Official Journal of the American Urological Association (AUA).

"For men undergoing [active surveillance](#), negative biopsies indicate low-volume disease and lower rates of disease progression," comments lead author Carissa E. Chu, MD, of University of California, San Francisco (UCSF) "These 'hidden' cancers have excellent long-term outcomes and remain ideal for continued active surveillance."

## 'Excellent' long-term outcomes with negative biopsies on active surveillance

During active surveillance, prostate cancer is carefully monitored for signs of progression through regular prostate-specific antigen (PSA) screening, prostate exams, imaging, and repeat biopsies. The goal is for men to avoid or delay treatment-related side effects without compromising such long-term outcomes as cancer progression or survival.

Sometimes, men undergoing active surveillance have negative biopsies showing no evidence of prostate cancer. While these patients may believe their cancer has "vanished," they most likely have low-volume or limited areas of prostate cancer that were not detected in the [biopsy](#) sample. Dr. Chu comments, "While a negative biopsy is good news, the long-term implications associated with such 'hidden' cancers remain unclear."

To evaluate the long-term significance of negative biopsies, the researchers analyzed 514 men

undergoing active surveillance for early-stage prostate cancer at UCSF between 2000 and 2019. All patients had at least three surveillance biopsies *after* their initial prostate cancer diagnosis (total four biopsies). Median follow-up time was nearly ten years.

Thirty-seven percent of patients had at least one negative biopsy during active surveillance, including 15 percent with consecutive negative biopsies. Men with negative surveillance biopsies had more favorable clinical characteristics, including low PSA density and fewer samples showing cancer at the initial prostate biopsy.

Negative biopsies were also associated with good long-term outcomes. At 10 years, rates of survival with no need for prostate cancer treatment (such as surgery or radiation) were 84 percent for men with consecutive negative biopsies, 74 percent for those with one negative biopsy, and 66 percent for those with no negative biopsies. After adjustment for other factors, men with one or more negative biopsies were much less likely to have cancer detected on a later biopsy.

However, having negative biopsies didn't mean that the cancer had "vanished"—even some men with consecutive negative biopsies later had positive biopsies or were diagnosed with a higher stage of cancer. Therefore surveillance, although less intense, is still recommended rather than "watchful waiting" for men in good health. Higher PSA density and suspicious findings on [prostate magnetic resonance](#) imaging (MRI) scans were associated with a higher risk of [cancer](#) detected on later biopsies.

"Negative surveillance biopsies in men on active surveillance indicate low-volume [prostate cancer](#) with very favorable outcomes," Dr. Chu and

colleagues conclude. "A less-intensive [surveillance](#) regimen should be supported in these patients after discussion of risks and benefits, particularly in those with low PSA density and adequate MRI-targeted sampling."

**More information:** Carissa E. Chu et al. The Clinical Significance of Multiple Negative Surveillance Prostate Biopsies for Men on Active Surveillance—Does Cancer Vanish or Simply Hide?, *Journal of Urology* (2020). DOI: [10.1097/JU.0000000000001339](https://doi.org/10.1097/JU.0000000000001339)

Provided by Wolters Kluwer Health

APA citation: 'Vanished' or 'hidden' prostate cancer? Men with negative biopsies during active surveillance have good outcomes (2020, November 17) retrieved 17 September 2022 from <https://medicalxpress.com/news/2020-11-hidden-prostate-cancer-men-negative.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*