


New study debunks blood type diet

December 4 2020



Credit: CC0 Public Domain

A study published in the *Journal of the Academy of Nutrition and Dietetics* by researchers with the Physicians Committee for Responsible Medicine—a nonprofit of 12,000 doctors—debunks the 'blood type diet' by finding that blood type was not associated with the effects of a plant-based diet on body weight, body fat, plasma lipid concentrations, or glycemic control.

This new study is based on a randomized control trial whose main findings were published in *JAMA Network Open* on Nov. 30. That trial randomly assigned overweight participants with no history of diabetes to an intervention or [control group](#) on a 1:1 ratio for 16 weeks. Participants in the intervention group followed a low-fat, plant-based diet. The control group made no diet changes. The key finding is that a plant-based diet ramps up metabolism as measured by an increase in after-meal calorie burn of 18.7%, on average, for the intervention group over the control.

To consider a potential connection between [blood type](#) and diet, researchers took the additional step of conducting a secondary analysis among intervention-group participants of the 16-week randomized clinical trial. They considered whether the effects of a plant-based dietary intervention on [body weight](#), [blood lipids](#), and [glycemic control](#) are associated with ABO blood type. The 'blood type diet' recommends a mainly plant-based diet for those with blood type A, while it recommends a diet heavy in meat for people with blood type O.

"We found that blood type made no difference," says study author Neal Barnard, MD, president of the Physicians Committee. "While the [blood type diet](#) says that a plant-based diet should be better for blood type A and less so for blood type O, it turned out to be beneficial for people of all blood types, and there was no evidence that meaty diets are good for anyone.

"Our research shows that all blood types benefit equally from a [vegan diet](#) based on the consumption of fruits and vegetables, legumes and whole grains, looking specifically at [weight loss](#) and cardiometabolic health in overweight adults," he says.

Main outcomes that were measured were body weight, fat mass, visceral fat volume, blood lipids, fasting plasma glucose, and HbA1c. T-tests

compared participants with blood type A to all other participants (non-A), and individuals with blood type O to all other participants (non-O).

There were no significant differences in any outcome between individuals of blood type A and non-A, or between individuals of blood type O and non-O. Mean body weight change was -5.7 kg for blood type A participants and -7.0 kg for non-A participants, and was -7.1 kg for type O participants and -6.2 kg for non-O participants. Mean total cholesterol decreased 17.2 mg/dl in the type A group and 18.3 mg/dl for non-A participants, and decreased 17.4 mg/dl among type O participants and 18.4 mg/dl for non-O participants.

More information: Neal D. Barnard et al, Blood Type Is Not Associated with Changes in Cardiometabolic Outcomes in Response to a Plant-Based Dietary Intervention, *Journal of the Academy of Nutrition and Dietetics* (2020). [DOI: 10.1016/j.jand.2020.08.079](https://doi.org/10.1016/j.jand.2020.08.079)

Provided by Physicians Committee for Responsible Medicine

Citation: New study debunks blood type diet (2020, December 4) retrieved 13 February 2024 from <https://medicalxpress.com/news/2020-12-debunks-blood-diet.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.