

# Common drug for build-up of blood following head injury worse than placebo, study finds

December 16 2020

---



Credit: Unsplash/CC0 Public Domain

A commonly-used treatment for chronic subdural haematoma—the build-up of 'old' blood in the space between the brain and the skull, usually as a

result of minor head injury—could lead to a worse outcome than receiving no medication, suggests new research from the University of Cambridge.

Chronic subdural haematoma is one of the most common neurological disorders and mainly affects older people. People affected often have headaches, deteriorating memory, confusion, balance problems or limb weakness. Surgery to drain the liquid collection is effective with the majority of patients improving.

A commonly used steroid, dexamethasone, has been used alongside surgery or instead of it since the 1970s. However, consensus has been lacking regarding the use of dexamethasone, especially since no high-quality studies confirming its effectiveness had been conducted until now.

With funding from the UK National Institute Health Research, a group of doctors and researchers from 23 neurosurgical units in the United Kingdom enrolled 748 patients with chronic subdural haematoma in the "Dexamethasone in Chronic Subdural Haematoma (Dex-CSDH)" randomized trial. A total of 375 patients were randomized to receive a two-week tapering course of dexamethasone and were compared with 373 patients randomized to an identical matching placebo.

The results of the study, published today in the *New England Journal of Medicine*, show that patients who received dexamethasone had a lower chance of favourable recovery at six months compared to patients who received placebo. More specifically, the vast majority of patients in both groups had an operation to drain the haematoma and had experienced significant functional improvement at six months compared to their initial admission to hospital.

Fewer patients in the dexamethasone group required repeat surgery for a

recurrent haematoma compared to patients in the placebo group. However, 84% of patients who received dexamethasone had recovered well at 6 months compared to 90% of patients who received placebo.

Peter Hutchinson, Professor of Neurosurgery at the University of Cambridge and the trial's Chief Investigator, said: "Chronic subdural haematoma has been steadily increasing in frequency over the past decades. Patients affected are often frail and have other co-existing medical conditions. Since the 1970s, dexamethasone has been used as a drug alongside or instead of surgery with a few studies reporting good results.

"Our trial sought to determine if dexamethasone should be offered routinely to all patients with chronic subdural haematoma or if its use should be abandoned. Based on our findings, we believe that dexamethasone should not be used in patients with chronic subdural haematoma anymore."

Angelos Kolias, Lecturer of Neurosurgery at the University of Cambridge and the trial's Co-chief Investigator, added: "The results of the study were surprising given that [dexamethasone](#) seemed to help reduce the number of repeat surgeries. However, this simply reinforces the importance of conducting high-quality trials with patient-reported outcomes as the main outcomes of interest".

Ellie Edlmann, the trial's research fellow, currently a Clinical Lecturer at the University of Plymouth, concluded: "Credit is due to all doctors and researchers from across the NHS who worked tirelessly in order to enrol all eligible patients in the trial; in particular, the role of trainee neurosurgeons, members of the British Neurosurgical Trainee Research Collaborative, needs to be highlighted. We sincerely thank all patients and their carers, as without their altruistic participation, this trial would not have been possible."

The trial was funded by the National Institute for Health Research (NIHR), with further support from the NIHR Cambridge Biomedical Research Centre, the NIHR Brain Injury MedTech Co-operative, the Royal College of Surgeons of England, and the Rosetrees Trust.

**More information:** *New England Journal of Medicine* (2020). [DOI: 10.1056/NEJMoa2020473](https://doi.org/10.1056/NEJMoa2020473)

Provided by University of Cambridge

Citation: Common drug for build-up of blood following head injury worse than placebo, study finds (2020, December 16) retrieved 30 December 2022 from <https://medicalxpress.com/news/2020-12-dexamethasone-patients-chronic-subdural-haematoma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.