

AASM: Multicomponent CBT recommended for chronic insomnia

23 December 2020



as a single-component therapy for treatment of chronic insomnia disorder.

"Although [sleep hygiene](#) practices are often suggested and well understood by patients, sleep hygiene [recommendations] do not constitute an effective stand-alone therapy," Edinger said in a statement.

Several authors disclosed financial ties to the pharmaceutical and medical technology industries.

More information: [Abstract/Full Text](#)
([subscription or payment may be required](#))

Copyright © 2020 [HealthDay](#). All rights reserved.

(HealthDay)—Multicomponent cognitive behavioral therapy is strongly recommended for the treatment of chronic insomnia in adults, according to a new clinical practice guideline issued by the American Academy of Sleep Medicine and published online Nov. 9 in the *Journal of Clinical Sleep Medicine*.

Jack D. Edinger, Ph.D., from National Jewish Health in Denver, and colleagues conducted a systematic literature review to evaluate the relevant literature and quality of evidence to develop recommendations for behavioral and psychological treatments for chronic insomnia disorder in adults.

The authors strongly recommend that clinicians use multicomponent [cognitive behavioral therapy](#) for treatment of chronic insomnia disorder in adults. Conditional recommendations include use of stimulus control, sleep restriction [therapy](#), and relaxation therapy as single-component therapies for treatment of chronic insomnia disorder in adults. Sleep hygiene is not recommended for use

APA citation: AASM: Multicomponent CBT recommended for chronic insomnia (2020, December 23) retrieved 15 May 2021 from <https://medicalxpress.com/news/2020-12-aasm-multicomponent-cbt-chronic-insomnia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.