


Sex differences in death after stroke

23 December 2020



Kornstein, MD, Executive Director of the Virginia Commonwealth University Institute for Women's Health, Richmond, VA.

More information: Hoang T. Phan et al, Sex Differences in Causes of Death After Stroke: Evidence from a National, Prospective Registry, *Journal of Women's Health* (2020). [DOI: 10.1089/jwh.2020.8391](https://doi.org/10.1089/jwh.2020.8391)

Provided by Mary Ann Liebert, Inc

Credit: CC0 Public Domain

Women were 39% more likely to die by 1 year after a first stroke. The sex difference was due to advanced age and more severe strokes in women, according to a new study in the *Journal of Women's Health*.

Among women and men with a first-ever stroke, women were approximately 7 years older. In addition, 9.3% fewer women could walk independently on admission to the hospital, suggestive of a more [severe stroke](#).

"Among those deceased by any cause, men had more deaths due to cancer (12% vs women 6%) and [ischemic heart disease](#) (8% vs women 6%) while women had more deaths attributed to stroke (50% vs men 41%) or other [cardiovascular disease](#) (16% vs men 13%), state Dominique Cadilhac, Ph.D., School of Clinical Sciences at Monash Health, and coauthors.

"Cadilhac and colleagues showed that women had a 65% greater risk of death associated with stroke. Not only were [women](#) more likely to be older at first stroke and to have greater stroke severity, but they were also less likely to be treated with aspirin for secondary stroke prevention," says *Journal of Women's Health* Editor-in-Chief Susan G.

APA citation: Sex differences in death after stroke (2020, December 23) retrieved 20 October 2022 from <https://medicalxpress.com/news/2020-12-sex-differences-death.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.