

Meeting India's air quality targets across south Asia may prevent 7% of pregnancy losses, modelling study estimates

January 7 2021



Credit: Unsplash/CC0 Public Domain

Poor air quality is associated with a considerable proportion of pregnancy loss in India, Pakistan, and Bangladesh, according to a modelling study published in *The Lancet Planetary Health* journal.

Previous studies have suggested a link between air pollution and [pregnancy loss](#) in other regions, but this is the first study to quantify the burden in south Asia, which is the most populous region in the world and has the highest rate of [pregnancy](#) loss. Therefore, understanding the [risk factors](#) for pregnancy loss in south Asia is crucial to improving maternal health regionally and globally.

Lead author on the study, Dr. Tao Xue, Peking University, China, says, "South Asia has the highest burden of pregnancy loss globally and is one of the most PM2.5 polluted regions in the world. Our findings suggest that [poor air quality](#) could be responsible for a considerable burden of pregnancy loss in the region, providing further justification for urgent action to tackle dangerous levels of pollution."

One of the co-authors, Dr. Tianjia Guan, is from the Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China. She says "We know losing a pregnancy can have knock-on mental, physical and economic effects on women, including increased risk of postnatal depressive disorders, infant mortality during subsequent pregnancy, and increase the costs related to pregnancy, such as loss of labour. Therefore, reducing pregnancy loss may also lead to knock-on improvements in gender equality."

To carry out their analysis, the authors combined data from household surveys on health from 1998-2016 (from women who reported at least one pregnancy loss and one or more livebirths) and estimated exposure to PM2.5 during pregnancy through combining satellite with atmospheric modelling outputs. They created a model to examine how exposure to PM2.5 increased women's risk of pregnancy loss, calculating risk for each $10 \mu\text{g}/\text{m}^3$ increased in PM2.5 after adjusting for maternal age, temperature and humidity, seasonal variation, and long-term trends in pregnancy loss.

Using this association, they calculated the number of pregnancy losses that may have been caused by PM_{2.5} in the whole region for the period 2000-16 and looked at how many pregnancy losses might have been prevented under India's and WHO's air quality standard (40 $\mu\text{g}/\text{m}^3$ and 10 $\mu\text{g}/\text{m}^3$, respectively).

In the study, they included 34,197 women who had lost a pregnancy, including 27,480 miscarriages and 6,717 stillbirths, which were compared to livebirth controls. Of the pregnancy loss cases, 77% were from India, 12% from Pakistan, and 11% from Bangladesh.

Gestational exposure to PM_{2.5} was associated with an increased likelihood of pregnancy loss, and this remained significant after adjusting for other factors. Each increase in 10 $\mu\text{g}/\text{m}^3$ was estimated to increase a mother's risk of pregnancy loss by 3%.

The increase in risk was greater for mothers from rural areas or those who became pregnant at an older age, compared to younger mothers from urban areas.

From 2000 to 2016, 349,681 pregnancy losses per year were associated with ambient exposure to air pollution exceeding India's air quality standard— accounting for 7% of the total annual pregnancy loss burden in this region. For air pollution above WHO air quality guideline, exposure may have contributed to 29% of pregnancy losses.

Although WHO's guidelines aims for a safer level of air pollution, the authors note that India's standard is a more realistic target level, given the high average levels of air pollution in the region and the need to balance practical governance and public health.

Pregnancy loss associated with [air pollution](#) was more common in the Northern plains region in India and Pakistan. Although the total burden

of pregnancy loss was predominantly borne by rural women aged under 30 years old in recent years, the burden attributable to PM2.5 also affected older mothers (aged 30 years or over) in [rural areas](#) because of their high susceptibility to the adverse effects of PM2.5.

The authors note several limitations of their study. In the surveys, they were not able to distinguish between natural pregnancy loss and abortions and there was under-reporting of pregnancy losses because of stigma or ignoring very early pregnancy losses. They also note that the survey data is subject to recall bias, therefore, recommending the causality of the association should be further examined in longitudinal studies. In addition, satellite-based estimates of PM2.5 were used, this was necessary because insufficient local monitoring is available.

More information: Estimation of pregnancy losses attributable to exposure to ambient fine particles in south Asia: an epidemiological case-control study, *The Lancet Planetary Health*, [DOI: 10.1016/S2542-5196\(20\)30268-0](#) , [www.thelancet.com/journals/lan ...](http://www.thelancet.com/journals/lan...)
[\(20\)30268-0/fulltext](#)

Provided by Lancet

Citation: Meeting India's air quality targets across south Asia may prevent 7% of pregnancy losses, modelling study estimates (2021, January 7) retrieved 28 January 2023 from <https://medicalxpress.com/news/2021-01-india-air-quality-south-asia.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--