

COVID-19 ups complication risks during childbirth

21 January 2021, by Steven Reinberg



Among the COVID-19 patients, the researchers found the relative risk of developing any type of blood clot was nearly five times higher than for those without the virus, and nearly four times higher for venous thromboembolism, clots in the veins.

These women were also far more likely to need [intensive care](#) or a ventilator, the researchers found.

Those who had the virus were:

- 7% more likely to need a C-section.
- 19% more likely to have preterm labor.
- 17% more likely to have a preterm delivery.
- 21% more likely to have preeclampsia.

There's little a [pregnant woman](#) can do to reduce these risks beyond not being infected, Jering said.

Women who have COVID-19 during childbirth are more likely to face complications than moms-to-be without the coronavirus, researchers say.

Fortunately, the absolute risk for complications for any one woman is very low (less than 1%). But the relative risks for problems—such as clotting and early labor—are significant, the new study found.

Still, "the findings here, truly, are that among [women](#) who are hospitalized for childbirth and who were diagnosed with COVID, adverse events are incredibly low. That should provide a lot of reassurance to women who are hoping to become pregnant during this period, or who are pregnant," said study co-author Dr. Karola Jering, from the cardiovascular medicine division at Brigham and Women's Hospital in Boston.

Over eight months in 2020, she and her colleagues collected data on more than 400,000 mothers-to-be, nearly 6,400 of whom were infected with COVID-19.

"The problem, of course, is that right now we mostly have supportive care for patients who have COVID, in general. And of the things that have been tested for treatment of patients with COVID, most of them have not been tested in pregnant women," said co-author Dr. Scott Solomon, also from Brigham and Women's.

But Jering said pregnant women are given the other drugs often given to COVID-19 patients, including blood thinners to prevent clots.

In sum, the study findings were positive, Jering stressed. Among the [pregnant women](#) with COVID-19 who gave birth, 99% were discharged home, 3% needed intensive care and 1% needed mechanical ventilation. Less than 1% died in the hospital.

Jering said that these findings should reassure women who have COVID-19 that, although complications can occur, most women will have a normal pregnancy and delivery.

Dr. Eran Bornstein is vice-chair of obstetrics and gynecology at Lenox Hill Hospital in New York City. He said, "Overall, these findings are important. They provide further support to prior observations regarding risk factors for COVID-19 during pregnancy as well as for pregnancy complications."

As previous research has shown, Hispanic and Black mothers were at greater risk of having the coronavirus, Bornstein noted. Young age, diabetes and obesity were also [risk factors](#).

"This is important, as it emphasizes the impact sociodemographic factors and [health conditions](#) have on the likelihood of having COVID-19 in pregnancy," Bornstein said.

The report was published online Jan. 15 in *JAMA Internal Medicine*.

More information: [Abstract/Full Text](#)

For more on pregnancy and COVID-19, see the [U.S. Centers for Disease Control and Prevention](#).

Copyright © 2020 [HealthDay](#). All rights reserved.

APA citation: COVID-19 ups complication risks during childbirth (2021, January 21) retrieved 20 September 2022 from <https://medicalxpress.com/news/2021-01-covid-ups-complication-childbirth.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.