


# Can't remember last night? 48% of drinkers have had a blackout by age 19

February 4 2021, by Wing See Yuen and Amy Peacock

---



Credit: AI-generated image ([disclaimer](#))

Alcohol-related blackouts aren't good for anybody's health, but they are particularly dangerous for young people.

Our [recent research](#) found blackouts are common once [young people](#) start drinking. At age 14, nearly one in ten adolescents who drank

alcohol in the past year had a [blackout](#).

By age 19, around 48% had experienced a blackout.

We also found around 14% of young Australians in our study had more and more alcohol-related blackouts as they aged through adolescence. Females were three times more likely to experience this increasing number of blackouts than males.

Young people who had this increasing number of blackouts were around 2.5 times more likely to develop severe alcohol problems including alcohol abuse and dependence in early adulthood.

To conduct this research, we [recruited](#) 1,821 13-year-olds from year 7 classes in New South Wales, Western Australia, and Tasmania in 2010-11. We asked them to complete surveys each year about their alcohol use.

We used eight years of data to analyze when they started drinking alcohol, if they had alcohol-related blackouts, and if they reported more severe alcohol harms, like [alcohol abuse](#) and dependence.

## **What is a blackout?**

An [alcohol-related blackout](#) happens when someone has [blood alcohol concentration](#) of about 0.15 or higher (three times the legal driving limit). Blackouts are more likely to be triggered when someone raises their [blood alcohol levels](#) very quickly, for example by "chugging" drinks or drinking on an empty stomach.

Despite the name, someone who's having an alcohol blackout is not unconscious (although people might become unconscious during or after a blackout). They can continue to do things such as talking and walking,

but afterwards they can't remember what they did while they were drunk. In other words, alcohol can temporarily stop your brain from forming long-term memories.

Most blackouts tend to be "spotty," where the person might be able to remember some things that happened while drinking but not others. With more severe blackouts, they aren't able to remember anything at all from when they started to black out, even if someone tries to remind them what happened.

## **Why are blackouts so dangerous?**

Alcohol affects everyone differently, so the number of drinks it takes to trigger a blackout varies from person to person.

Regardless of the number of drinks consumed, when someone has a blackout, it means they're drinking at a level that affects their memory and behavior.

Because young peoples' brains are still developing until they're about 25, they're very [vulnerable](#) to the damage alcohol [causes to the brain](#). Drinking regularly at amounts leading to blackouts can cause [permanent brain damage](#).

In the short term, someone having a blackout has reached blood-alcohol levels that make them more likely to undertake [risky activities](#) such as driving, having [unprotected sex](#), and other behaviors that can lead to [injury](#) or death.

Blackouts are also linked to having [problems](#) with work, school and social life.

In the long term, young adults who have alcohol blackouts are 1.6-2.6

times as likely to experience alcohol-related injuries [two years later](#) and around 1.5 times as likely to have symptoms of alcohol dependence [five years later](#).

## Who's particularly at risk?

Having lower bodyweight, drinking faster, and not eating before drinking all [increase](#) the [chances](#) of having a blackout.

Females are also around 1.8 times [more likely](#) to have a blackout when drinking the [same amount](#) as males. This is because females are, on average, smaller than males and have less water in their bodies to dilute consumed alcohol, so they absorb alcohol into their blood faster than males.

## How can you prevent blackouts?

If you're going to drink alcohol, these tips can help prevent blackouts:

- make sure to eat before and during a drinking session
- try to sip your drink rather than taking gulps or chugging
- have some water between each alcoholic drink
- avoid binge drinking (having four or more drinks in two hours).

### [One 2018 study showed](#)

young people tend to be aware that drinking alcohol on an empty stomach leads to blackouts. But less than one in four of the participants were aware that females are at greater risk because of how their body processes alcohol.

Teachers can help by teaching young people about the factors that

increase their chances of having a blackout.

Parents can also play an important role in helping their children learn to have a healthier relationship with alcohol by doing things like [reducing their own binge drinking](#) and not [supplying alcohol](#) to anyone under the age of 18.

Regardless of your age, it's never too late to rethink your relationship with [alcohol](#). If you've had a blackout, it's a very good indicator you're [drinking](#) at a concerning level, regardless of how many drinks led to the blackout.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Can't remember last night? 48% of drinkers have had a blackout by age 19 (2021, February 4) retrieved 4 July 2024 from <https://medicalxpress.com/news/2021-02-night-drinkers-blackout-age.html>

|  |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|