

Commuters are inhaling unacceptably high levels of carcinogens

February 15 2021



Driver exercising extreme caution in his car. Credit: Stan Lim/UCR

A new study finds that California's commuters are likely inhaling chemicals at levels that increase the risk for cancer and birth defects.

As with most chemicals, the poison is in the amount. Under a certain threshold of exposure, even known carcinogens are not likely to cause

cancer. Once you cross that threshold, the risk for disease increases.

Governmental agencies tend to regulate that threshold in workplaces. However, private spaces such as the interior of our cars and living rooms are less studied and less regulated.

Benzene and formaldehyde—both used in automobile manufacturing—are known to cause cancer at or above certain levels of exposure and are Prop. 65-listed chemicals.

New UC Riverside research shows that the average commuter in California is exceeding the threshold for exposure, breathing in unsustainably high levels of both chemicals.

Both benzene and formaldehyde are carcinogens, and benzene carries the additional risk of reproductive and developmental toxicity.

"These chemicals are very volatile, moving easily from plastics and textiles to the air that you breathe," said David Volz, UCR professor of environmental toxicology.

The study, published in the journal *Environment International*, calculated the daily dose of [benzene](#) and formaldehyde being inhaled by drivers with commutes of at least 20 minutes per day.

It found that up to 90% of the population in Los Angeles, San Diego, Orange, Santa Clara, and Alameda counties have at least a 10% chance of exceeding cancer risk from inhaling the chemicals, based on having 30-minute average commute times.

"Of course, there is a range of exposure that depends on how long you're in the car, and how much of the compounds your car is emitting," said Aalekhya Reddam, a [graduate student](#) in the Volz laboratory, and lead

author of the study.

Previously, Volz and Reddam [studied](#) commuter exposure to a flame retardant called TDCIPP or chlorinated tris, and found that longer commute times increased exposure to that carcinogen as well.

They set out on this study wanting to understand the risk of that compound relative to other chemicals introduced during car manufacturing.

Reddam advises commuters to keep the windows open during their rides if possible. "At least with some air flow, you'd be diluting the concentration of these chemicals inside your car," she said.

Benzene is used to produce synthetic fibers, and formaldehyde is a binder in plastics. "There should be alternatives to these chemicals to achieve the same goals during vehicle manufacturing," Volz said. "If so, these should be used."

More information: Aalekhya Reddam et al, Inhalation of two Prop 65-listed chemicals within vehicles may be associated with increased cancer risk, *Environment International* (2021). [DOI: 10.1016/j.envint.2021.106402](#)

Provided by University of California - Riverside

Citation: Commuters are inhaling unacceptably high levels of carcinogens (2021, February 15) retrieved 9 July 2023 from <https://medicalxpress.com/news/2021-02-commuters-inhaling-unacceptably-high-carcinogens.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.