

Are there differences in the brains of autistic men and women?

12 March 2021



Credit: CC0 Public Domain

Around three times as many males are diagnosed with autism than females. This suggests that biological sex factors may play a role in the development and presentation of autism. Studies on the neurobiology (brain biology) of males and females with autism have begun to examine brain networks but results have been mixed. This is largely due to the limited availability of data from autistic females.

In response, researchers from Child Mind Institute and colleagues involved in the AIMS2TRIALS, have combined thousands of MRI data openly available for scientific discovery in the Autism Brain Imaging Exchange (ABIDE) repository to explore [brain](#) network differences between autistic and neurotypical control males and females. They used the ABIDE sample for discovery of new information and two additional large samples to see if those findings could be repeated (i.e., replicated). These included one sample derived from the Gender Explorations of Neurogenetics and Development to Advance Autism Research made available through the National Database for Autism Research and another one shared by the collaborators of the

AIMS2TRIALS.

Across these three samples, the researchers found that both neurotypical males and autistic people showed reduced resting-state brain function in the so-called 'default network', a network that is active when we engage in social cognition or thoughts about ourselves. Additionally, in the discovery [sample](#) and in one of the largest of the two replication samples, it was shown that connections between hemispheres (the two halves of the brain) in the visual cortex are reduced in autistic [females](#), while autistic males are not different from [males](#) who are not autistic. The results suggest that many autistic people may have different interactions between the two hemispheres of their brain when compared to non-autistic people. This reflects a combination of effects, including some that appear to be unrelated to sex, and some in which there is an interaction between sex and autism diagnosis. Each of these effects appears specific to a different system in the brain.

This study highlights the importance of data sharing and collaboration for implementing discovery science and addressing critical challenges related to reproducibility of findings—which affect all of fields of science. The researchers suggest that there remains an urgent need for more research with similarly large groups of participants, as only then do studies have enough statistical power to reliably account for sources of variability and therefore generate robust conclusions. Until now, limited replication of imaging findings has hampered brain imaging research in [autism](#). The open sharing policies of the Autism Brain Imaging Data Exchange and the NIMH Data Archive, through which the Gender Explorations of Neurogenetics Development to Advance Autism Research was made available, are particularly promising for accelerating the pace of advancement.

More information: Dorothea L. Floris et al, Towards robust and replicable sex differences in

the intrinsic brain function of autism, *Molecular Autism* (2021). DOI: [10.1186/s13229-021-00415-z](https://doi.org/10.1186/s13229-021-00415-z)

Provided by The Child Mind Institute

APA citation: Are there differences in the brains of autistic men and women? (2021, March 12) retrieved 3 June 2022 from <https://medicalxpress.com/news/2021-03-differences-brains-autistic-men-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.